Index of Authors

ADIGA, P. R. *see* MUNIYAPPA, K. 887–894

AFTING, E.-G., BECKER, M. L. & ELCE, J. S. Proteinase and proteinase-inhibitor activities of rat uterine myometrium during pregnancy and involution 99–106

AL-AYASH, A. I. & WILSON, M. T. The mechanism of reduction of single-site redox proteins by ascorbic acid 641–648

ALDRIGE, W. N. *see* ELLIOTT, B. M. 461–470

ALHADEFF, J. A. & ANDREWS-SMITH, G. L. Subunit composition of human liver α-L-fucosidase 753–756

AMBLER, R. P., DANIEL, M., MEYER, T. E., BARTSCH, R. G. & KAMEN, M. D. The amino acid sequence of cytochrome *c'* from the purple sulphur bacterium *Chromatium vinosum* 819–823

ANDREWS-SMITH, G. L. *see* ALHADEFF, J. A. 753–756

ASHANI, Y. & LEADER, H. Formation of an unstable covalent intermediate during the inhibition of electric-eel acetylcholinesterase with 1,3,2-dioxaphosphorinane 2-oxides 781–790

ATGER, M., MERCIER, J.-C., HAZE, G., FRIDLANSKY, F. & MILGROM, E. *N*-Terminal sequences of uteroglobin and its precursor 985–988

AUGUSTUS, B. W. & HUTCHINSON, D. W. The synthesis of phenyl[2-^{3}H]glyoxal 377–379

AWASTHI, Y. C., DAO, D. D., LAL, A. K. & SRIVASTAVA, S. K. Purification and properties of glutathione peroxidase from human placenta 471–476

AZZI, A. *see* SIGRIST-NELSON, K. 687–692

BADEN, H. P. *see* LEE, L. D. 187–196

BAINES, B. S. & BROCKLEHURST, K. A necessary modification to the preparation of papain from any high-quality latex of *Carica papaya* and evidence for the structural integrity of the enzyme produced by traditional methods 541–548

BAKER, T. S., JENNISON, K. M. & KELLIE, A. E. The direct radioimmunoassay of oestrogen glucuronides in human female urine 729–738

BANNISTER, J. V. *see* CASS, A. E. G. 477–486

BANNISTER, W. H. *see* CASS, A. E. G. 477–486

BARBER, D. *see* WALSH, T. A. 29–39

BARKER, R., BODEN, N., CAYLEY, G., CHARLTON, S. C., HENSON, R., HOLMES, M. C., KELLY, I. D. & KNOWLES, P. F. Properties of cupric ions in benzylamine oxidase from pig plasma as studied by magnetic-resonance and kinetic methods 289–302

BARTSCH, R. G. *see* AMBLER, R. P. 819–823

BARUA, A. B., VERMA, K. & DAS, S. R. Structure and synthesis of naturally occurring anhydrovitamin A_2 791–799

BATTE, E. A. *see* MIAN, N. 319–330

BAYLEY, P. M. *see* HOOD, K. 425–432

BECKER, M. L. *see* AFTING, E.-G. 99–106

BEELEY, J. G. *see* FRANÇOIS-GERARD, C. 679–685

BÉGIN-HEICK, N. *see* TONG, J. H. 943–949

BEIS, I. *see* SCRUTTON, M. C. 833–846

BEN-YOSEPH, Y. *see* FIDDLER, M. B. 175–180

BERKHOUT, T. A. *see* PASTINK, A. 917–923

BERLET, H. H. *see* LEHNERT, T. 813–818

BLOXHAM, D. P., SHARMA, R. P. & WILTON, D. C. A detailed investigation of the properties of lactate dehydrogenase in which the 'essential' cysteine-165 is modified by thioalkylation 769–780

BODEN, N., HOLMES, M. C. & KNOWLES, P. F. Properties of the cupric sites in bovine superoxide dismutase studied by nuclear-magnetic-relaxation measurements 303–309

BODEN, N. *see also* BARKER, R. 289–302

BOOTH, N. A., CAMPBELL, R. D. & FOTHERGILL, J. E. The purification and characterization of bovine C4, the fourth component of complement 959–965

BOOTH, N. A. *see also* CAMPBELL, R. D. 531–540

BOUCEK, R. J., NOBLE, N. L. & GUNJA-SMITH, Z. A possible role for dehydroxylysinonorleucine in collagen fibre and bundle formation 853–860

BRAMHALL, J. S., SHIFLETT, M. A. & WISNIESKI, B. J. Mapping the membrane proteins of Newcastle-disease virus with a photoreactive glycolipid probe 765–768

BRATCHER, S. C. *see* NITTA, K. 385–392

BRAY, R. C., GUTTERIDGE, S., STOTTER, D. A. & TANNER, S. J. The mechanism of action of xanthine oxidase. The relationship between the Rapid and Very Rapid molybdenum electron-paramagnetic-resonance signals 357–360

BRETT, M. & FINDLAY, J. B. C. Investigation of the organization of rhodopsin in the sheep photoreceptor membrane by using cross-linking reagents 215–223

BRIDGES, J. W. *see* ELLIOTT, B. M. 461–470

BROCKLEHURST, K. *see* BAINES, B. S. 541–548

BRYANT, T. H. E. *see* MACKIE, C. 977–979

BUCKINGHAM, K. W. *see* HENG-KHOO, C. S. 559–567

BURCHELL, B. *see* LALANI, E.-N. M. A. 993–995

CAMPBELL, R. D., BOOTH, N. A. & FOTHERGILL, J. E. Purification and characterization of subcomponent C1q of the first component of bovine complement 531–540

CAMPBELL, R. D. *see also* BOOTH, N. A. 959–965

CAMPOS-CAVIERES, M. *see* ORLACCHIO, A. 583–593

CARNE, T., TIPPING, E. & KETTERER, B. The binding and catalytic activities of forms of ligandin after modification of its thiol groups 433–439

CASEY, R. Immunoaffinity chromatography as a means of purifying legumin from *Pisum* (pea) seeds 509–520

CASS, A. E. G., HILL, H. A. O., BANNISTER, J. V. & BANNISTER, W. H. Zinc(II) binding to apo-(bovine erythrocyte superoxide dismutase) 477–486

CAYLEY, G. *see* BARKER, R. 289–302

CHAMBERS, S. M. *see* RENWICK, A. G. C. 401–408

CHANCE, B., SARONIO, C. & LEIGH, J. S., JR. Compound C_2, a product of the reaction of oxygen and the mixed-valence state of cytochrome oxidase. Optical evidence for a type-1 copper 931–941

CHANCE, E. M. *see* CLORE, G. M. 613–621

CHARLTON, S. C. *see* BARKER, R. 289–302

CHAUDHURI, S. & THOMAS, E. W. Affinity chromatography of pig heart fumarase 115–119

CHRAMBACH, A. *see* NEWBY, A. C. 623–630

CLARKE, C. M. & HARTLEY, B. S. Purification, properties and specificity of the restriction endonuclease from *Bacillus stearothermophilus* 49–62

CLORE, G. M. & CHANCE, E. M. Low-temperature kinetics of the reaction of fully reduced membrane-bound cytochrome oxidase with oxygen in the Soret, α and near-infrared regions 613–621

CLOTHIER, B. & JOHNSON, M. K. Rapid aging of neurotoxic esterase after inhibition by di-isopropyl phosphorofluoridate 549–558
COHEN-SOLAL, L., LIAN, J. B., KOSSIVA, D. & GLIMCHER, M. J. Identification of organic phosphorus covalently bound to collagen and non-collagenous proteins of chicken-bone matrix. The presence of *O*-phosphoserine and *O*-phosphothreonine in non-collagenous proteins, and their absence from phosphorylated collagen 81–98
COLBY, J. & DALTON, H. Characterization of the second prosthetic group of the flavoenzyme NADH–acceptor reductase (component C) of the methane monooxygenase from *Methylococcus capsulatus* (Bath) 903–908
COLBY, J. *see also* STIRLING, D. I. 361–364
COOPER, C. S. & QUINCEY, R. V. Convolution analysis of transcription by yeast DNA-dependent ribonucleic acid polymerase A. A mathematical method for studying ribonucleic acid chain elongation 825–831
CORFIELD, A. P., SCHAUER, R. & WEMBER, M. The preparation of CMP-sialic acids by using CMP-acylneuraminate synthase from frog liver immobilized on Sepharose 4B 1–7
COWEN, D. M. *see* MIAN, N. 319–330
COX, D. *see* TSITILOU, S. G. 331–346
CRANG, A. J. & RUMSBY, M. G. The intrinsic fluorescence of isolated central-nervous-system myelin-sheath preparations 739–745
CRAVEN, A. H. *see* TOPPING, R. M. 501–508
CRONLUND, A. L. *see* KAGAN, H. M. 203–214
CRUTCHER, S. E. & GEARY, P. J. Properties of the iron–sulphur proteins of the benzene dioxygenase system from *Pseudomonas putida* 393–400
CZEKANSKI, S. *see* POZNANSKY, M. J. 989–991

DADAY, A., LAMBERT, G. R. & SMITH, G. D. Measurement *in vivo* of hydrogenase-catalysed hydrogen evolution in the presence of nitrogenase enzyme in cyanobacteria 139–144
DALTON, H. *see* COLBY, J. 903–908; STIRLING, D. I. 361–364
DANIEL, M. *see* AMBLER, R. P. 819–823
DAO, D. D. *see* AWASTHI, Y. C. 471–476
DAS, S. R. *see* BARUA, A. B. 791–799
DAVIDSON, W. S. & FLYNN, T. G. Kinetics and mechanism of action of aldehyde reductase from pig kidney 595–601
DAVIES, R. C. & NEUBERGER, A. Control of 5-aminolaevulinate synthetase activity in *Rhodopseudomonas spheroides*. Binding of pyridoxal phosphate to 5-aminolaevulinate synthetase 661–671
DAVIES, R. C. & NEUBERGER, A. Control of 5-aminolaevulinate synthetase activity in *Rhodopseudomonas spheroides*. Purification and properties of the high-activity form of the enzyme 649–659
DICKINSON, F. M. *see* ELECTRICWALA, A. H. 449–459
DIONYSSIOU-ASTERIOU, A. & RAKITZIS, E. T. Activation of cathepsin D by glycine ethyl ester 355–356
D'IORIO, A. *see* TONG, J. H. 943–949
DIXON, H. B. F. Derivation of molecular p*K* values from pH-dependences 249–250
DOUMENG, C. & MAROUX, S. Aminotripeptidase, a cytosol enzyme from rabbit intestinal mucosa 801–808
DOYEN, N. & LAPRESLE, C. Partial non-cleavage by cyanogen bromide of a methionine–cystine bond from human serum albumin and bovine α-lactalbumin 251–254
DRYSDALE, G. R. *see* WEIDMAN, S. W. 169–174
DUŠEK, Z. *see* POHLREICH, P. 707–719

EGMOND, M. R. *see* WHITEHEAD, E. P. 631–639
ELAHI, E. & SELLINGER, O. Z. The postnatal methylation of transfer ribonucleic acid in brain. Evidence for the methylation of precursor transfer ribonucleic acid 381–384
ELCE, J. S. *see* AFTING, E.-G. 99–106
ELECTRICWALA, A. H. & DICKINSON, F. M. Kinetic studies on dogfish liver glutamate dehydrogenase 449–459
ELLIOTT, B. M., ALDRIGE, W. N. & BRIDGES, J. W. Triethyltin binding to cat haemoglobin. Evidence for two chemically distinct sites and a role for both histidine and cysteine residues 461–470
EL-YASSIN, D. I., FELL, D. A., LLOYD, B. B. & FISHER, R. B. The breakdown of 2,3-bis(phospho)-D-glycerate by Fe(III)-haemoglobin 373–375

FELL, D. A. *see* EL-YASSIN, D. I. 373–375
FIDDLER, M. B., BEN-YOSEPH, Y. & NADLER, H. L. Binding of human liver hydrolases by immobilized lectins 175–180
FIEDLER, F. *see* LEMON, M. 159–168
FINCH, S. A. E., SLATER, T. F. & STIER, A. Nucleotide metabolism by microsomal UDP-glucuronyltransferase and nucleoside diphosphatase as determined by ^{31}P nuclear-magnetic-resonance spectroscopy 925–930
FINDLAY, J. B. C. *see* BRETT, M. 215–223
FISHER, R. B. *see* EL-YASSIN, D. I. 373–375
FITTON, J. E. & SHAW, W. V. Comparison of chloramphenicol acetyltransferase variants in staphylococci. Purification, inhibitor studies and *N*-terminal sequences 575–582
FLANAGAN, P. R. & FORSTNER, G. G. Enzyme activity in partly dissociated fragments of rat intestinal maltase/glucoamylase 487–492
FLYNN, T. G. *see* DAVIDSON, W. S. 595–601
FÖRG-BREY, B. *see* LEMON, M. 159–168
FORSTNER, G. G. *see* FLANAGAN, P. R. 487–492
FOTHERGILL, J. E. *see* BOOTH, N. A. 959–965; CAMPBELL, R. D. 531–540
FOWLER, L. J. *see* JOHN, R. A. 721–728
FRANÇOIS-GERARD, C., GERDAY, C. & BEELEY, J. G. Turtle-dove ovomucoid, a glycoprotein proteinase inhibitor with P_1-blood-group antigen activity 679–685
FREEZE, H., GEIGER, B. & MILLER, A. L. Carbohydrate composition of human placental *N*-acetylhexosaminidase A and B 749–752
FRÈRE, J.-M. *see* MARQUET, A. 909–916
FRIDLANSKY, F. *see* ATGER, M. 985–988
FRITZ, H. *see* LEMON, M. 159–168

GACESA, P. & VENN, R. F. The preparation of stable enzyme–coenzyme complexes with endogenous catalytic activity 369–372
GAUSH, C. R. *see* HOUGLAND, A. E. 311–317
GAZZOLO, L. *see* QUASH, G. 275–282
GEARY, P. J. *see* CRUTCHER, S. E. 393–400

GEIGER, B. *see* FREEZE, H. 749–752
GERDAY, C. *see* FRANÇOIS-GERARD, C. 679–685
GHAZARIAN, J. G. *see* KULKOSKI, J. A. 673–678
GHUYSEN, J.-M. *see* MARQUET, A. 909–916
GIBBONS, G. F. & PULLINGER, C. R. Utilization of endogenous and exogenous sources of substrate for cholesterol biosynthesis by isolated hepatocytes 255–263
GLIMCHER, M. J. *see* COHEN-SOLAL, L. 81–98
GÓRNICKI, S. Z. *see* HAMILTON, T. A. 197–201
GRAND, R. J. A., PERRY, S. V. & WEEKS, R. A. Troponin C-like proteins (calmodulins) from mammalian smooth muscle and other tissues 521–529
GREEN, N. M. *see* KNIGHT, C. G. 225–236
GREENWOOD, C. *see* WALSH, T. A. 29–39
GRIGAT, K.-P., KOPPE, K., SEUFERT, C.-D. & SÖLING, H.-D. Acetyl-coenzyme A deacylase activity in liver is not an artifact. Subcellular distribution and substrate specificity of acetyl-coenzyme A deacylase activities in rat liver 71–79
GUNJA-SMITH, Z. *see* BOUCEK, R. J. 853–860
GUTTERIDGE, S. *see* BRAY, R. C. 357–360

HALE, G. & PERHAM, R. N. Polypeptide-chain stoicheiometry and lipoic acid content of the pyruvate dehydrogenase complex of *Escherichia coli* 129–137
HAMES, B. D. & HODSON, B. A. Accumulation of UDP-glucose pyrophosphorylase by axenically grown amoebae of *Dictyostelium discoideum* 21–28
HAMILTON, T. A., GÓRNICKI, S. Z. & SUSSMAN, H. H. Alkaline phosphatase from human milk. Comparison with isoenzymes from placenta and liver 197–201
HARDINGHAM, T. E. The role of link-protein in the structure of cartilage proteoglycan aggregates 237–247
HARTLEY, B. S. *see* CLARKE, C. M. 49–62
HATLEN, L. *see* HEGENAUER, J. 693–695
HAYASHI, T. T. & MACFARLANE, K. Comparison of endogenous and exogenous RNA primers of poly(U) polymerase in rat hepatic ribosomes 895–902
HAZE, G. *see* ATGER, M. 985–988
HEGENAUER, J., SALTMAN, P. & HATLEN, L. Removal of cadmium(II) from crystallized ferritin 693–695
HENG-KHOO, C. S., RUCKER, R. B. & BUCKINGHAM, K. W. Additional evidence for a proform to tropoelastin from chick aorta 559–567
HENSON, R. *see* BARKER, R. 289–302
HERRIES, D. G. *see* MIAN, N. 319–330
HILL, H. A. O. *see* CASS, A. E. G. 477–486
HIRSCHAUER, C. *see* LEMON, M. 159–168
HISKEY, R. G. *see* SCOTT, M. E. 879–886
HODSON, B. A. *see* HAMES, B. D. 21–28
HOLMES, M. C. *see* BARKER, R. 289–302; BODEN, N. 303–309
HOOD, K., BAYLEY, P. M. & ROBERTS, G. C. K. Circular-dichroism studies of ligand binding to dihydrofolate reductase from *Lactobacillus casei* MTX/R 425–432
HOUGLAND, A. E., GAUSH, C. R. & MAYBERRY, W. R. Glycolipids and fatty acids of two dog kidney cell lines 311–317
HRADEC, J. *see* POHLREICH, P. 707–719
HUTCHINSON, D. W. *see* AUGUSTUS, B. W. 377–379

ISERN DE CALDENTEY, M. & WHEELER, K. P. Requirement for negatively charged dispersions of phospholipids for interaction with lipid-depleted adenosine triphosphatase 265–273

JAKOBSON, I., WARHOLM, M. & MANNERVIK, B. Multiple inhibition of glutathione *S*-transferase A from rat liver by glutathione derivatives: kinetic analysis supporting a steady-state random sequential mechanism 861–868
JÄNNE, J. *see* KALLIO, A. 63–69
JENNISON, K. M. *see* BAKER, T. S. 729–738
JOHN, R. A., JONES, E. D. & FOWLER, L. J. Enzyme-induced inactivation of transaminases by acetylenic and vinyl analogues of 4-aminobutyrate 721–728
JOHN, R. A. *see also* WILLIAMS, J. A. 121–127
JOHNSON, M. K. *see* CLOTHIER, B. 549–558; WALSH, T. A. 29–39
JONES, E. D. *see* JOHN, R. A. 721–728

KAGAN, H. M., SULLIVAN, K. A., OLSSON, T. A., III & CRONLUND, A. L. Purification and properties of four species of lysyl oxidase from bovine aorta 203–214
KALLIO, A., LÖFMAN, M., PÖSÖ, H. & JÄNNE, J. Diamine-induced inhibition of liver ornithine decarboxylase 63–69
KAMEN, M. D. *see* AMBLER, R. P. 819–823
KELLIE, A. E. *see* BAKER, T. S. 729–738
KELLY, I. D. *see* BARKER, R. 289–302
KEOLOUANGKHOT, T. *see* QUASH, G. 275–282
KETTERER, B. *see* CARNE, T. 433–439
KNIGHT, C. G. & GREEN, N. M. Interaction of dinitrophenyl groups bound to bovine serum albumin with univalent fragments of anti-dinitrophenyl antibody 225–236
KNOTT-HUNZIKER, V., ORLEK, B. S., SAMMES, P. G. & WALEY, S. G. 6β-Bromopenicillanic acid inactivates β-lactamase I 365–367
KNOWLES, P. F. *see* BARKER, R. 289–302; BODEN, N. 303–309
KOEHLER, K. A. *see* SCOTT, M. E. 879–886
KOPPE, K. *see* GRIGAT, K.-P. 71–79
KOSSIVA, D. *see* COHEN-SOLAL, L. 81–98
KRESS, B. C. & MILLER, A. L. Urinary lysosomal hydrolases in mucolipidosis II and mucolipidosis III 409–415
KŘIŽ, O. *see* POHLREICH, P. 707–719
KRONMAN, M. J. *see* NITTA, K. 385–392
KUAN, K. N., LEE, Y. Y. & MELIUS, P. An improved method of light-induced pigmentation 981–983
KUBILUS, J. *see* LEE, L. D. 187–196
KULKOSKI, J. A. & GHAZARIAN, J. G. Purification and characterization of the ferredoxin component of 25-hydroxycholecalciferol 1α-hydroxylase 673–678

LAL, A. K. *see* AWASTHI, Y. C. 471–476
LALANI, E.-N. M. A. & BURCHELL, B. Stimulation of defective Gunn-rat liver uridine diphosphate glucuronyltransferase activity *in vitro* by alkyl ketones 993–995
LAMBERT, G. R. *see* DADAY, A. 139–144
LAPRESLE, C. *see* DOYEN, N. 251–254
LEADER, H. *see* ASHANI, Y. 781–790
LEE, L. D., KUBILUS, J. & BADEN, H. P. Intraspecies heterogeneity of epidermal keratins isolated from bovine hoof and snout 187–196
LEE, Y. Y. *see* KUAN, K. N. 981–983
LEHNERT, T. & BERLET, H. H. Selective inactivation of lactate dehydrogenase of rat tissues by sodium deoxycholate 813–818

LEIGH, J. S., JR. *see* CHANCE, B. 931–941
LEMON, M., FIEDLER, F., FÖRG-BREY, B., HIRSCHAUER, C., LEYSATH, G. & FRITZ, H. The isolation and properties of pig submandibular kallikrein 159–168
LEYSATH, G. *see* LEMON, M. 159–168
LIAN, J. B. *see* COHEN-SOLAL, L. 81–98
LIMSON-ZAMORA, M. *see* TONG, J. H. 943–949
LLOYD, B. B. *see* EL-YASSIN, D. I. 373–375
LOEFFLER, R. S. T., SINNOTT, M. L., SYKES, B. D. & WITHERS, S. G. Interaction of the *lacZ* β-galactosidase of *Escherichia coli* with some β-D-galactopyranoside competitive inhibitors 145–152
LOFFET, A. *see* MARQUET, A. 909–916
LÖFMAN, M. *see* KALLIO, A. 63–69
LYNN, W. S. *see* SAHU, S. 153–158

MACFARLANE, K. *see* HAYASHI, T. T. 895–902
MACKIE, C., BRYANT, T. H. E. & MOWBRAY, J. A rapid and sensitive method for measuring ribonucleic acid in ribosomal preparations 977–979
MAGER, W. H. *see* PASTINK, A. 917–923
MANNERVIK, B. *see* JAKOBSON, I. 861–868; UOTILA, L. 869–878
MARGISON, G. P., MARGISON, J. M. & MONTESANO, R. Persistence of methylated bases in ribonucleic acid of Syrian golden hamster liver after administration of dimethylnitrosamine 967–973
MARGISON, J. M. *see* MARGISON, G. P. 967–973
MAROUX, S. *see* DOUMENG, C. 801–808
MARQUET, A., FRÈRE, J.-M., GHUYSEN, J.-M. & LOFFET, A. Effects of nucleophiles on the breakdown of the benzylpenicilloyl–enzyme complex EI* formed between benzylpenicillin and the exocellular DD-carboxypeptidase–transpeptidase of *Streptomyces* strain R61 909–916
MATHIAS, A. P. *see* TSITILOU, S. G. 331–346
MAYBERRY, W. R. *see* HOUGLAND, A. E. 311–317
MCGURK, K. S. & SPIVEY, H. O. Ligand-induced conformational transitions and secondary structure composition of chicken liver pyruvate carboxylase 697–705
MELIUS, P. *see* KUAN, K. N. 981–983
MERCIER, J.-C. *see* ATGER, M. 985–988
MEYER, T. E. *see* AMBLER, R. P. 819–823
MIAN, N., HERRIES, D. G., COWEN, D. M. & BATTE, E. A. The multiple forms and kinetic properties of the *N*-acetyl-β-D-hexosaminidases from colonic tumours and mucosa of rats treated with 1,2-dimethylhydrazine 319–330
MILGROM, E. *see* ATGER, M. 985–988
MILLER, A. L. *see* FREEZE, H. 749–752; KRESS, B. C. 409–415
MOAK, G. S. *see* WAPNIR, R. A. 347–352
MONTESANO, R. *see* MARGISON, G. P. 967–973
MORELLO, A. & REPETTO, Y. UDP-glucosyltransferase activity of housefly microsomal fraction 809–812
MORRISSEY, P. G. *see* SHREVE, M. R. 761–763
MOWBRAY, J. *see* MACKIE, C. 977–979
MUNIYAPPA, K. & ADIGA, P. R. Isolation and characterization of thiamin-binding protein from chicken egg white 887–894
MUNN, E. A. *see* ORLACCHIO, A. 583–593

NADLER, H. L. *see* FIDDLER, M. B. 175–180
NEUBERGER, A. *see* DAVIES, R. C. 649–659, 661–671
NEWBY, A. C. & CHRAMBACH, A. Disaggregation of adenylate cyclase during polyacrylamide-gel electrophoresis in mixtures of ionic and non-ionic detergents 623–630
NIMMO, H. G. Evidence for the existence of isoenzymes of glycerol phosphate acyltransferase 283–288
NITTA, K., BRATCHER, S. C. & KRONMAN, M. J. Anomalous reaction of 4-chloro-7-nitrobenzofurazan with thiol compounds 385–392
NOBLE, N. L. *see* BOUCEK, R. J. 853–860
NOTARIO, V. *see* VILLA, T. G. 107–114

O'BRIEN, P. J. *see* SHREVE, M. R. 761–763
OLSSON, T. A., III *see* KAGAN, H. M. 203–214
ONON, E. O. Purification and partial characterization of the exotoxin of *Corynebacterium ovis* 181–186
ORLACCHIO, A., CAMPOS-CAVIERES, M., PASHEV, I. & MUNN, E. A. Some kinetic and other properties of the isoenzymes of aspartate aminotransferase isolated from sheep liver 583–593
ORLEK, B. S. *see* KNOTT-HUNZIKER, V. 365–367

PASHEV, I. *see* ORLACCHIO, A. 583–593
PASTINK, A., BERKHOUT, T. A., MAGER, W. H. & PLANTA, R. J. Analysis of histones from the yeast *Saccharomyces carlsbergensis* 917–923
PERHAM, R. N. *see* HALE, G. 129–137
PERRY, S. V. *see* GRAND, R. J. A. 521–529
PLANTA, R. J. *see* PASTINK, A. 917–923
POHLREICH, P., KŘIŽ, O., TUHÁČKOVÁ, Z., DUŠEK, Z. & HRADEC, J. Purification and some properties of a protein factor binding and deacylating initiator transfer ribonucleic acid 707–719
PÖSÖ, H. *see* KALLIO, A. 63–69
POZNANSKY, M. J. & CZEKANSKI, S. Cholesterol exchange as a function of cholesterol/phospholipid mole ratios 989–991
PRICE, N. C. The reaction of rabbit muscle creatine kinase with some derivatives of iodoacetamide 603–612
PULLINGER, C. R. *see* GIBBONS, G. F. 255–263

QUASH, G., KEOLOUANGKHOT, T., GAZZOLO, L., RIPOLL, H. & SAEZ, S. Diamine oxidase and polyamine oxidase activities in normal and transformed cells 275–282
QUINCEY, R. V. *see* COOPER, C. S. 825–831

RAKITZIS, E. T. *see* DIONYSSIOU-ASTERIOU, A. 355–356
RENWICK, A. G. C., CHAMBERS, S. M. & WILLCOX, P. 'Affinity' chromatography of steroid-transforming enzymes with a non-steroidal ligand 401–408
REPETTO, Y. *see* MORELLO, A. 809–812
RIDGE, D. *see* TSITILOU, S. G. 331–346
RIDING, G. A. *see* WILLADSEN, P. 41–47
RIPOLL, H. *see* QUASH, G. 275–282
ROBERTS, G. C. K. *see* HOOD, K. 425–432
RUCKER, R. B. *see* HENG-KHOO, C. S. 559–567
RUMSBY, M. G. *see* CRANG, A. J. 739–745

SAEZ, S. *see* QUASH, G. 275–282
SAHU, S. & LYNN, W. S. Characterization of a high-molecular-weight glycoprotein isolated from the pulmonary secretions of patients with alveolar proteinosis 153–158
SALTMAN, P. *see* HEGENAUER, J. 693–695
SAMMES, P. G. *see* KNOTT-HUNZIKER, V. 365–367

SANDY, J. D. The assay of xylosyltransferase in cartilage extracts. A modified procedure for preparation of Smith-degraded proteoglycan 569–574
SARONIO, C. *see* CHANCE, B. 931–941
SCHAUER, R. *see* CORFIELD, A. P. 1–7
SCOTT, M. E., KOEHLER, K. A. & HISKEY, R. G. The effects of calcium ions and pH on bovine prothrombin fragment 1. Intrinsic fluorescence studies 879–886
SCRUTTON, M. C. & BEIS, I. Inhibitory effects of histidine and their reversal. The roles of pyruvate carboxylase and N^{10}-formyltetrahydrofolate dehydrogenase 833–846
SEILMAN, S. *see* TOPPING, R. M. 493–499
SELLINGER, O. Z. *see* ELAHI, E. 381–384
SEUFERT, C.-D. *see* GRIGAT, K.-P. 71–79
SHARMA, R. P. *see* BLOXHAM, D. P. 769–780
SHAW, W. V. *see* FITTON, J. E. 575–582
SHIFLETT, M. A. *see* BRAMHALL, J. S. 765–768
SHINKAI, H. & YONEMASU, K. Hydroxylysine-linked glycosides of human complement subcomponent C1q and of various collagens 847–852
SHREVE, M. R., MORRISSEY, P. G. & O'BRIEN, P. J. Lipid and steroid hydroperoxides as substrates for the non-selenium-dependent glutathione peroxidase 761–763
SIGRIST-NELSON, K. & AZZI, A. Purification of the chloroplast-membrane dicyclohexylcarbodi-imide-binding proteolipid by ion-exchange chromatography 687–692
SIMMONDS, R. J. *see* YON, R. J. 417–424
SINNOTT, M. L. *see* LOEFFLER, R. S. T. 145–152
SLATER, T. F. *see* FINCH, S. A. E. 925–930
SMITH, G. D. *see* DADAY, A. 139–144
SÖLING, H.-D. *see* GRIGAT, K.-P. 71–79
SPIVEY, H. O. *see* MCGURK, K. S. 697–705
SPRINGALL, J. P. *see* WALSH, T. A. 29–39
SRIVASTAVA, S. K. *see* AWASTHI, Y. C. 471–476
STIER, A. *see* FINCH, S. A. E. 925–930
STIRLING, D. I., COLBY, J. & DALTON, H. A comparison of the substrate and electron-donor specificities of the methane mono-oxygenases from three strains of methane-oxidizing bacteria 361–364
STOTTER, D. A. *see* BRAY, R. C. 357–360
SULLIVAN, K. A. *see* KAGAN, H. M. 203–214
SUNDARAM, T. K. *see* WRIGHT, I. P. 441–448
SUSSMAN, H. H. *see* HAMILTON, T. A. 197–201
SYKES, B. D. *see* LOEFFLER, R. S. T. 145–152

TANNER, S. J. *see* BRAY, R. C. 357–360
THOMAS, E. W. *see* CHAUDHURI, S. 115–119
THOMSON, A. J. *see* WALSH, T. A. 29–39
TIPPING, E. *see* CARNE, T. 433–439
TONG, J. H., LIMSON-ZAMORA, M., D'IORIO, A. & BÉGIN-HEICK, N. Liver monoamine oxidase in the obese–hyperglycaemic (*ob*/*ob*) mouse 943–949
TOPPING, R. M. & CRAVEN, A. H. The identification of distinctive forms of human α_2-macroglobulin by using the numerical relationship between trypsin binding in α- and β-modes 501–508
TOPPING, R. M. & SEILMAN, S. A four-straight-line model for the proteinase-binding characteristics of human blood serum 493–499
TSITILOU, S. G., COX, D., MATHIAS, A. P. & RIDGE, D. The characterization of the non-histone chromosomal proteins of the main classes of nuclei from rat brain fractionated by zonal centrifugation 331–346
TUHÁČKOVÁ, Z. *see* POHLREICH, P. 707–719

UMEZURIKE, G. M. The cellulolytic enzymes of *Botryodiplodia theobromae* Pat. Separation and characterization of cellulases and β-glucosidases 9–19
UOTILA, L. & MANNERVIK, B. A steady-state-kinetic model for formaldehyde dehydrogenase from human liver. A mechanism involving NAD^+ and the hemimercaptal adduct of glutathione and formaldehyde as substrates and free glutathione as an allosteric activator of the enzyme 869–878

VENN, R. F. *see* GACESA, P. 369–372
VERMA, K. *see* BARUA, A. B. 791–799
VILLA, T. G., NOTARIO, V. & VILLANUEVA, J. R. Occurrence of an endo-1,3-β-glucanase in culture fluids of the yeast *Candida utilis*. Purification and characterization of the enzyme activity 107–114
VILLANUEVA, J. R. *see* VILLA, T. G. 107–114
VINCENT, S. P. Oxidation–reduction potentials of molybdenum and iron–sulphur centres in nitrate reductase from *Escherichia coli* 757–759

WALEY, S. G. *see* KNOTT-HUNZIKER, V. 365–367
WALSH, T. A., JOHNSON, M. K., GREENWOOD, C., BARBER, D., SPRINGALL, J. P. & THOMSON, A. J. Some magnetic properties of *Pseudomonas* cytochrome oxidase 29–39
WAPNIR, R. A. & MOAK, G. S. β-2-Thienyl-DL-alanine as an inhibitor of phenylalanine hydroxylase and phenylalanine intestinal transport 347–352
WARHOLM, M. *see* JAKOBSON, I. 861–868
WEEKS, R. A. *see* GRAND, R. J. A. 521–529
WEIDMAN, S. W. & DRYSDALE, G. R. The biosynthesis of methylcitrate 169–174
WEMBER, M. *see* CORFIELD, A. P. 1–7
WHEELER, K. P. *see* ISERN DE CALDENTEY, M. 265–273
WHITEHEAD, E. P. & EGMOND, M. R. Patterns of apparent co-operativity in a simple random non-equilibrium enzyme–substrate–modifier mechanism. Comparison with equilibrium allosteric models 631–639
WILLADSEN, P. & RIDING, G. A. Characterization of a proteolytic-enzyme inhibitor with allergenic activity. Multiple functions of a parasite-derived protein 41–47
WILLCOX, P. *see* RENWICK, A. G. C. 401–408
WILLIAMS, J. A. & JOHN, R. A. Generation of aspartate aminotransferase multiple forms by deamidation 121–127
WILSON, M. T. *see* AL-AYASH, A. I. 641–648
WILTON, D. C. The mechanism of adduct formation between NAD^+ and pyruvate bound to pig heart lactate dehydrogenase 951–957
WILTON, D. C. *see also* BLOXHAM, D. P. 769–780
WISNIESKI, B. J. *see* BRAMHALL, J. S. 765–768
WITHERS, S. G. *see* LOEFFLER, R. S. T. 145–152
WRIGHT, I. P. & SUNDARAM, T. K. Simple efficient methods for the isolation of malate dehydrogenase from thermophilic and mesophilic bacteria 441–448

YON, R. J. & SIMMONDS, R. J. The adsorption of proteins and protein–dodecyl sulphate complexes on *N*-(3-carboxypropionyl)aminodecyl-Sepharose 417–424
YONEMASU, K. *see* SHINKAI, H. 847–852

Index of Subjects

[2-^{3}H]Acetophenone, synthesis of phenyl[2-^{3}H]glyoxal from, and its use in the modification of arginine residues in enzymes (Augustus, B. W. & Hutchinson, D. W.) 377–379

Acetylcholinesterase, electric-eel, formation of an unstable covalent intermediate during the inhibition by 1,3,2-dioxaphosphorinane 2-oxides of the activity of (Ashani, Y. & Leader, H.) 781–790

Acetyl-coenzyme A deacylase, subcellular distribution and substrate specificity of the activities of, in rat liver (Grigat, K.-P., Koppe, K., Seufert, C.-D. & Söling, H.-D.) 71–79

β-*N*-Acetylhexosaminidase, placental, human, determination of the carbohydrate compositions of the A and B isoenzymes of (Freeze, H., Geiger, B. & Miller, A. L.) 749–752

β-*N*-Acetylhexosaminidases, multiple forms and kinetic properties of, from colonic tumours and colonic mucosa from rats treated with 1,2-dimethylhydrazine (Mian, N., Herries, D. G., Cowen, D. M. & Batte, E. A.) 319–330

Acids, dibasic, derivation of molecular p*K* values of, from dependences on pH (Dixon, H. B. F.) 249–250

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, kidney, rabbit, lipid-depleted, requirement for negatively charged dispersions of phospholipids for interaction with (Isern de Caldentey, M. & Wheeler, K. P.) 265–273

Adenylate cyclase, plasma-membrane, liver, rat, disaggregation of, during polyacrylamide-gel electrophoresis in mixtures of ionic and non-ionic detergents (Newby, A. C. & Chrambach, A.) 623–630

Albumin, serum, human, partial non-cleavage by cyanogen bromide of a methionine–cystine bond in (Doyen, N. & Lapresle, C.) 251–254

Albumin, serum, ox, effects of sodium dodecyl sulphate on the adsorption of, and other proteins on *N*-(3-carboxypropionyl)aminodecyl-Sepharose (Yon, R. J. & Simmonds, R. J.) 417–424

Albumin, serum, ox, interaction of dinitrophenyl groups bound to, with univalent fragments of anti-dinitrophenyl antibody (Knight, C. G. & Green, N. M.) 225–236

Alcohol dehydrogenase, comparison of the kinetics and mechanism of action of pig kidney aldehyde reductase with those of (Davidson, W. S. & Flynn, T. G.) 595–601

Aldehyde reductase, kidney, pig, kinetics and mechanism of action of (Davidson, W. S. & Flynn, T. G.) 595–601

Alkaline phosphatase, *see* Phosphatase, alkaline

Alkyl ketones, stimulation *in vitro* by, of the activity of defective Gunn-rat liver uridine diphosphate glucuronyltransferase (Lalani, E.-N. M. A. & Burchell, B.) 993–995

Alveolar proteinosis, characterization of a high-molecular-weight glycoprotein isolated from the pulmonary secretions of human patients with (Sahu, S. & Lynn, W. S.) 153–158

Amino acid sequence, determination of, of *Chromatium vinosum* A.T.C.C. 17899 cytochrome *c'* (Ambler, R. P., Daniel, M., Meyer, T. E., Bartsch, R. G. & Kamen, M. D.) 819–823

Amino acid sequences, *N*-terminal, comparison of, of variants of *Staphylococcus* spp. chloramphenicol acetyltransferase (Fitton, J. E. & Shaw, W. V.) 575–582

Amino acid sequences, *N*-terminal, determination of, of rabbit uterus uteroglobulin and its precursor (Atger, M., Mercier, J.-C., Haze, G., Fridlansky, F. & Milgrom, E.) 985–988

4-Aminobutyrate aminotransferase, brain, rabbit, enzyme-induced inactivation by acetylenic and vinyl analogues of 4-aminobutyrate of, and other aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728

4-Aminobutyrate, enzyme-induced inactivation by acetylenic and vinyl analogues of, of various mammalian aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728

5-Aminolaevulinate synthetase, *Rhodopseudomonas spheroides* N.C.I.B. 8253, purification and properties of the high-activity form of (Davies, R. C. & Neuberger, A.) 649–659

5-Aminolaevulinate synthetase, role of the binding of pyridoxal 5′-phosphate in the control of the activity of, in *Rhodopseudomonas spheroides* N.C.I.B. 8253 (Davies, R. C. & Neuberger, A.) 661–671

Aminotripeptidase, cytosolic, mucosal, small-intestinal, rabbit, purification and properties of (Doumeng, C. & Maroux, S.) 801–808

Anabaena cylindrica, measurement *in vivo* of the evolution of hydrogen by, that is catalysed by hydrogenase independently of nitrogenase (Daday, A., Lambert, G. R. & Smith, G. D.) 139–144

Anhydrovitamin A_2, *Saccobranchus fossilis*, naturally occurring, identification of, as 3-hydroxyanhydroretinol (Barua, A. B., Verma, K. & Das, S. R.) 791–799

Antibody, anti-dinitrophenyl, interaction of dinitrophenyl groups bound to ox serum albumin with univalent fragments of (Knight, C. G. & Green, N. M.) 225–236

Anti-dinitrophenyl antibody, interaction of dinitrophenyl groups bound to ox serum albumin with univalent fragments of (Knight, C. G. & Green, N. M.) 225–236

Aorta, chick, additional evidence for the existence of a precursor to tropoelastin in (Heng-Khoo, C. S., Rucker, R. B. & Buckingham, K. W.) 559–567

Aorta, ox, purification and properties of four species of lysyl oxidase from (Kagan, H. M., Sullivan, K. A., Olsson, T. A., III & Cronlund, A. L.) 203–214

Apo-(superoxide dismutase), erythrocyte, ox, binding of zinc ions to (Cass, A. E. G., Hill, H. A. O., Bannister, J. V. & Bannister, W. H.) 477–486

Arginine residues, synthesis of phenyl[2-^{3}H]glyoxal from [2-^{3}H]acetophenone and its use in the modification of, in enzymes (Augustus, B. W. & Hutchinson, D. W.) 377–379

Ascorbate, mechanism of the reduction by, of single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648

Aspartate aminotransferase, heart, pig, enzyme-induced inactivation by acetylenic and vinyl analogues of 4-aminobutyrate of, and other aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728

Aspartate aminotransferase, heart, pig, generation by deamidation of multiple forms of (Williams, J. A. & John, R. A.) 121–127
Aspartate aminotransferase, mitochondrial, liver, sheep, purification of, and comparison of its kinetic and other properties with those of the cytosolic isoenzyme (Orlacchio, A., Campos-Cavieres, M., Pashev, I. & Munn, E. A.) 583–593
12-(4-Azido-2-nitrophenoxy)stearoylglucosamine, use of, as a photoreactive probe for the mapping of the envelope proteins of Newcastle-disease virus (Bramhall, J. S., Shiflett, M. A. & Wisnieski, B. J.) 765–768
Azurin, *Pseudomonas aeruginosa*, mechanism of the reduction by ascorbate of, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648

Bacillus caldotenax, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Bacillus cereus, inactivation by 6β-bromopenicillanic acid of β-lactamase I from (Knott-Hunziker, V., Orlek, B. S., Sammes, P. G. & Waley, S. G.) 365–367
Bacillus stearothermophilus N.C.I.B. 8924, purification, properties and specificity of restriction endodeoxyribonuclease from (Clarke, C. M. & Hartley, B. S.) 49–62
Bacillus stearothermophilus, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Bacillus subtilis, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Benzene dioxygenase system, *Pseudomonas putida*, properties of the iron–sulphur proteins of (Crutcher, S. E. & Geary, P. J.) 393–400
Benzylamine oxidase, plasma, pig, study by magnetic-resonance and kinetic methods of the properties of cupric ions in (Barker, R., Boden, N., Cayley, G., Charlton, S. C., Henson, R., Holmes, M. C., Kelly, I. D. & Knowles, P. F.) 289–302
Benzylpenicillin, effects of nucleophiles on the breakdown of the benzylpenicilloyl–enzyme complex EI* formed between *Streptomyces* R61 exocellular DD-carboxypeptidase–peptidyltransferase and (Marquet, A., Frère, J.-M., Ghuysen, J.-M. & Loffet, A.) 909–916
2,3-Bis(phospho)-D-glycerate, breakdown of, catalysed by human ferrihaemoglobin (El-Yassin, D. I., Fell, D. A., Lloyd, B. B. & Fisher, R. B.) 373–375
Blood plasma, pig, study by magnetic-resonance and kinetic methods of the properties of cupric ions in benzylamine oxidase from (Barker, R., Boden, N., Cayley, G., Charlton, S. C., Henson, R., Holmes, M. C., Kelly, I. D. & Knowles, P. F.) 289–302
Blood serum, human, four-straight-line model for the characteristics of the binding of ox trypsin by (Tipping, R. M. & Seilman, S.) 493–499
Blood-group antigen, P_1, human, purification of turtle-dove egg-white ovomucoid and its activity as a proteinase inhibitor and as (François-Gerard, C., Gerday, C. & Beeley, J. G.) 679–685
Bombyx mori, *see* Silkworm
Bone, chicken, presence of *O*-phosphoserine and *O*-phosphothreonine residues in non-collagenous proteins and their absence from phosphorylated collagen from (Cohen-Solal, L., Lian, J. B., Kossiva, D. & Glimcher, M. J.) 81–98
Boophilus microplus, *see* Tick
Botryodiplodia theobromae, separation and characterization of cellulases and β-glucosidases of (Umezurike, G. M.) 9–19
Brain, chicken, rapid aging of neurotoxic esterase from, after inhibition by di-isopropyl phosphorofluoridate (Clothier, B. & Johnson, M. K.) 549–558
Brain, ox, intrinsic fluorescence of preparations of myelin sheath isolated from (Crang, A. J. & Rumsby, M. G.) 739–745
Brain, rabbit, enzyme-induced inactivation by acetylenic and vinyl analogues of 4-aminobutyrate of 4-aminobutyrate aminotransferase from, and other aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728
Brain, rat, characterization of the chromosomal non-histone proteins of the main classes of nuclei from, fractionated by zonal centrifugation (Tsitilou, S. G., Cox, D., Mathias, A. P. & Ridge, D.) 331–346
Brain, rat, evidence for the methylation of the precursors of transfer ribonucleic acids in, during postnatal development (Elahi, E. & Sellinger, O. Z.) 381–384
Brain, rat, selective inactivation by sodium deoxycholate of lactate dehydrogenases from, and other tissues (Lehnert, T. & Berlet, H. H.) 813–818
6β-Bromopenicillanic acid, inactivation by, of *Bacillus cereus* lactamase I (Knott-Hunziker, V., Orlek, B. S., Sammes, P. G. & Waley, S. G.) 365–367

Cadmium(II), removal of, from crystallized horse spleen ferritin (Hegenauer, J., Saltman, P. & Halten, L.) 693–695
Calcium ions, effects of pH and, on the intrinsic fluorescence of ox prothrombin fragment 1 (Scott, M. E., Koehler, K. A. & Hiskey, R. G.) 879–886
Calmodulins, purification and characterization of, from rabbit smooth muscle and other tissues (Grand, R. J. A., Perry, S. V. & Weeks, R. A.) 521–529
Candida utilis C.E.C.T. 1061, purification and characterization of endo-1,3-β-glucanase secreted into the culture medium by (Villa, T. G., Notario, V. & Villanueva, J. R.) 107–114
Carbohydrate compositions, determination of, of the A and B isoenzymes of human placental β-*N*-acetylhexosaminidase (Freeze, H., Geiger, B. & Miller, A. L.) 749–752
DD-Carboxypeptidase–peptidyltransferase, exocellular, *Streptomyces* R61, effects of nucleophiles on the breakdown of the benzylpenicilloyl–enzyme complex EI* formed between benzylpenicillin and (Marquet, A., Frère, J.-M., Ghuysen, J.-M. & Loffet, A.) 909–916
N-(3-Carboxypropionyl)aminodecyl-Sepharose, effects of sodium dodecyl sulphate on the adsorption of proteins on (Yon, R. J. & Simmonds, R. J.) 417–424
Carica papaya, *see* Papaya
Cartilage, chick-embryo, modified procedure for the preparation of Smith-degraded proteoglycans for use as substrates in the assay of the activity of xylosyltransferase in extracts of (Sandy, J. D.) 569–574

Cartilage, laryngeal, pig, role of link-protein in the structure of aggregates of hyaluronate and proteoglycans from (Hardingham, T. E.) 237–247
Cathepsin D, splcen, ox, activation by glycine ethyl ester of (Dionyssiou-Asteriou, A. & Rakitzis, E. T.) 355–356
Cellulases, *Botryodiplodia theobromae*, separation and characterization of, and β-glucosidases (Umezurike, G. M.) 9–19
Central nervous system, *see* Nervous system, central
Chick, additional evidence for the existence of a precursor to tropoelastin in aorta of (Heng-Khoo, C. S., Rucker, R. B. & Buckingham, K. W.) 559–567
Chick embryo, modified procedure for the preparation of Smith-degraded proteoglycans for use as substrates in the assay of the activity of xylosyltransferase in extracts of cartilage of (Sandy, J. D.) 569–574
Chick, purification and characterization of the ferredoxin component of kidney 25-hydroxycholecalciferol 1α-hydroxylase from (Kulkoski, J. A. & Ghazarian, J. G.) 673–678
Chicken, 'affinity' chromatography of liver oestradiol-17β dehydrogenase from, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408
Chicken egg-white, isolation and characterizatic : of thiamin-binding protein from, and its interaction with riboflavin-binding protein (Muniyappa, K. & Adiga, P. R.) 887–894
Chicken, ligand-induced conformational transitions and secondary structure composition of liver pyruvate carboxylase from (McGurk, K. S. & Spivey, H. O.) 697–705
Chicken, presence of *O*-phosphoserine and *O*-phosphothreonine residues in non-collagenous proteins and their absence from phosphorylated collagen from bone of (Cohen-Solal, L., Lian, J. B., Kossiva, D. & Glimcher, M. J.) 81–98
Chicken, rapid aging of neurotoxic esterase from brain of, after inhibition by di-isopropyl phosphorofluoridate (Clothier, B. & Johnson, M. K.) 549–558
Chloramphenicol acetyltransferase, *Staphylococcus* spp., purification, inhibitor studies and *N*-terminal amino acid sequences of variants of (Fitton, J. E. & Shaw, W. V.) 575–582)
4-Chloro-7-nitrobenzofurazan, anomalous reaction of, with thiols (Nitta, K., Bratcher, S. C. & Kronman, M. J.) 385–392
4-Chloro-7-nitro-2-oxa-1,3-diazole, *see* 4-Chloro-7-nitrobenzofurazan
Chloroplasts, lettuce-leaf, purification by ion-exchange chromatography of the membrane dicyclohexylcarbodiimide-binding proteolipid from (Sigrist-Nelson, K. & Azzi, A.) 687–692
Cholecalciferol, 25-hydroxy-, purification and characterization of the ferredoxin component of the chick kidney enzyme responsible for the 1α-hydroxylation of (Kulkoski, J. A. & Ghazarian, J. G.) 673–678
Cholesterol, effect of the ratio of, to phospholipid in human erythrocyte-'ghost' membrane on cholesterol exchange (Poznansky, M. J. & Czekanski, S.) 989–991
Cholesterol, utilization of endogenous and exogenous sources of substrate for the biosynthesis of, by isolated rat hepatocytes (Gibbons, G. F. & Pullinger, C. R.) 255–263
Chromatin, *Saccharomyces carlsbergensis*, isolation and characterization of histones from (Pastink, A., Berkhout, T. A., Mager, W. H. & Planta, R. J.) 917–923
Chromatium vinosum A.T.C.C. 17899, determination of the amino acid sequence of cytochrome *c'* from (Ambler, R. P., Daniel, M., Meyer, T. E., Bartsch, R. G. & Kamen, M. D.) 819–823
Chromosomes, characterization of the non-histone proteins of, of the main classes of rat brain nuclei fractionated by zonal centrifugation (Tsitilou, S. G., Cox, D., Mathias, A. P. & Ridge, D.) 331–346
α-Chymotrypsin, activation by, of silkworm tyrosinase precursor and its relevance to light-induced pigmentation (Kuan, K. N., Lee, Y. Y. & Melius, P.) 981–983
Citrate synthase, heart, pig, formation of diastereoisomers of methylcitrate catalysed by, and its relevance to the appearance of methylcitrate in the urine of human patients with propionic acidaemia (Weidman, S. W. & Drysdale, G. R.) 169–174
Collagen fibres and bundles, possible role for dehydrodihydroxylysinonorleucine in the formation of, in rat connective tissue (Boucek, R. J., Noble, N. L. & Gunja-Smith, Z.) 853–860
Collagen, phosphorylated, absence of *O*-phosphoserine and *O*-phosphothreonine residues from, and their presence in non-collagenous protein from chicken bone (Cohen-Solal, L., Lian, J. B., Kossiva, D. & Glimcher, M. J.) 81–98
Collagens, various, characterization of the hydroxylysine-linked glycosides of human complement subcomponent C1q and of (Shinkai, H. & Yonemasu, K.) 847–852
Colon, rat, multiple forms and kinetic properties of β-*N*-acetylhexosaminidases from mucosa of, and from colonic tumours from animals treated with 1,2-dimethylhydrazine (Mian, N., Herries, D. G., Cowen, D. M. & Batte, E. A.) 319–330
Complement, human, characterization of the hydroxylysine-linked glycosides of subcomponent C1q of, and of various collagens (Shinkai, H. & Yonemasu, K.) 847–852
Complement, ox, purification and characterization of component C4 from (Booth, N. A., Campbell, R. D. & Fothergill, J. E.) 959–965
Complement, ox, purification and characterization of subcomponent C1q from (Campbell, R. D., Booth, N. A. & Fothergill, J. E.) 531–540
Component C1, complement, human, characterization of the hydroxylysine-linked glycosides of subcomponent C1q of, and of various collagens (Shinkai, H. & Yonemasu, K.) 847–852
Component C1, complement, ox, purification and characterization of subcomponent C1q of (Campbell, R. D., Booth, N. A. & Fothergill, J. E.) 531–540
Component C4, complement, ox, purification and characterization of (Booth, N. A., Campbell, R. D. & Fothergill, J. E.) 959–965
Connective tissue, rat, possible role for dehydrodihydroxylysinonorleucine in the formation of collagen fibres and bundles in (Boucek, R. J., Noble, N. L. & Gunja-Smith, Z.) 853–860

Copper ion sites, study by nuclear-magnetic-relaxation measurements of the properties of, of ox erythrocyte superoxide dismutase (Boden, N., Holmes, M. C. & Knowles, P. F.) 303–309
Copper ions, study by magnetic-resonance and kinetic methods of the properties of, of pig plasma benzylamine oxidase (Barker, R., Boden, N., Cayley, G., Charlton, S. C., Henson, R., Holmes, M. C., Kelly, I. D. & Knowles, P. F.) 289–302
Copper, type-I, evidence for the presence of, in a product (Compound C_2) of the reaction of oxygen with the mixed-valence state of ox heart mitochondrial cytochrome *c* oxidase (Chance, B., Saronio, C. & Leigh, J. S., Jr.) 931–941
Cortisone reductase, *Streptomyces dehydrogenans*, 'affinity' chromatography of, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408
Corynebacterium ovis, purification and partial characterization of the extracellular toxin produced by (Onon, E. O.) 181–186
Coryphaena hippurus, *see* Dolphin fish
Creatine kinase, skeletal-muscle, rabbit, reaction of some derivatives of iodoacetamide with (Price, N. C.) 603–612
Cupric ion sites, study by nuclear-magnetic-relaxation measurements of the properties of, of ox erythrocyte superoxide dismutase (Boden, N., Holmes, M. C. & Knowles, P. F.) 303–309
Cupric ions, study by magnetic-resonance and kinetic methods of the properties of, of pig plasma benzylamine oxidase (Barker, R., Boden, N., Cayley, G., Charlton, S. C., Henson, R., Holmes, M. C., Kelly, I. D. & Knowles, P. F.) 289–302
Cysteine residue 165, 'essential', properties of pig heart lactate dehydrogenase in which, has been modified by thioalkylation (Bloxham, D. P., Sharma, R. P. & Wilton, D. C.) 769–780
Cysteine residues, evidence for two chemically distinct sites and a role for histidine residues and, in the binding of triethyltin to cat haemoglobin (Elliott, B. M., Aldridge, W. N. & Bridges, J. W.) 461–470
Cytidine monophosphate acylneuraminate synthase, liver, frog, synthesis of cytidine monophosphate sialic acids by using, immobilized on Sepharose 4B (Corfield, A. P., Schauer, R. & Wember, M.) 1–7
Cytidine monophosphate sialic acids, synthesis of, by using frog liver cytidine monophosphate acylneuraminate synthase immobilized on Sepharose 4B (Corfield, A. P., Schauer, R. & Wember, M.) 1–7
Cytochrome *c*, heart, horse, mechanism of the reduction by ascorbate of, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Cytochrome *c* oxidase, membrane-bound, mitochondrial, heart, ox, fully reduced, low-temperature kinetics of the reaction of, with oxygen studied by spectroscopy in the Soret, α-band and near-infrared regions (Clore, G. M. & Chance, E. M.) 613–621
Cytochrome *c* oxidase, mitochondrial, heart, ox, evidence for the presence of type-I copper in a product (Compound C_2) of the reaction of oxygen with the mixed-valence state of (Chance, B., Saronio, C. & Leigh, J. S., Jr.) 931–941
Cytochrome *c'*, *Chromatium vinosum* A.T.C.C. 17899, determination of the amino acid sequence of (Ambler, R. P., Daniel, M., Meyer, T. E., Bartsch, R. G. & Kamen, M. D.) 819–823
Cytochrome c_{551}, *Pseudomonas aeruginosa*, mechanism of the reduction by ascorbate of, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Cytochrome oxidase, *Pseudomonas aeruginosa* N.C.T.C. 6750, some magnetic properties of (Walsh, T. A., Johnson, M. K., Greenwood, C., Barber, D., Springall, J. P. & Thomson, A. J.) 29–39
Cytosol, liver, rat, purification and properties of a protein factor from, that binds and deacylates initiator transfer ribonucleic acid (Pohlreich, P., Kříž, O., Tuháčková, Z., Dušek, Z. & Hradec, J.) 707–719
Cytosol, mucosal, small-intestinal, rabbit, purification and properties of aminotripeptidase from (Doumeng, C. & Maroux, S.) 801–808

Dehydrodihydroxylysinonorleucine, possible role for, in the formation of collagen fibres and bundles in rat connective tissue (Boucek, R. J., Noble, N. L. & Gunja-Smith, Z.) 853–860
Deoxycholate, sodium, selective inactivation by, of lactate dehydrogenases from various rat tissues (Lehnert, T. & Berlet, H. H.) 813–818
Deoxyribonuclease, endo-, restriction, *Bacillus stearothermophilus* N.C.I.B. 8924, purification, properties and specificity of (Clarke, C. M. & Hartley, B. S.) 49–62
Deoxyribonucleic acid, use of convolution analysis as a mathematical method for the study of the transcription of, by yeast ribonucleic acid polymerase A (Cooper, C. S. & Quincey, R. V.) 825–831
Deoxyribonucleic acids, specificity of the cleavage of, by *Bacillus stearothermophilus* N.C.I.B. 8924 restriction endodeoxyribonuclease (Clarke, C. M. & Hartley, B. S.) 49–62
Development, postnatal, evidence for the methylation of the precursors of rat brain transfer ribonucleic acids during (Elahi, E. & Sellinger, O. Z.) 381–384
Diamine oxidase, activities of polyamine oxidase and, in normal and transformed rat kidney cells (Quash, G., Keolouangkhot, T., Gazzolo, L., Ripoll, H. & Saez, S.) 275–282
Diamines, induction by, of inhibition of the activity of ornithine decarboxylase in rat liver (Kallio, A., Löfman, M., Pösö, H. & Jänne, J.) 63–69
Dictyostelium discoideum, accumulation of uridine diphosphate glucose pyrophosphorylase by axenically grown amoebae of (Hames, B. D. & Hodson, B. A.) 21–28
Dictyostelium discoideum, mechanism of the reduction by ascorbate of cytochrome *c* from, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Dihydrofolate reductase, *see* Tetrahydrofolate dehydrogenase
Di-isopropyl phosphorofluoridate, rapid aging of chicken brain neurotoxic esterase after inhibition by (Clothier, B. & Johnson, M. K.) 549–558

1,2-Dimethylhydrazine, multiple forms and kinetic properties of β-*N*-acetylhexosaminidases from colonic tumours and colonic mucosa from rats treated with (Mian, N., Herries, D. G., Cowen, D. M. & Batte, E. A.) 319–330
Dimethylnitrosamine, persistence of methylated base residues in hamster liver ribonucleic acids after administration of, to the animals (Margison, G. P., Margison, J. M. & Montesano, R.) 967–973
Dinitrophenyl groups, interaction of, bound to ox serum albumin with univalent fragments of anti-dinitrophenyl antibody (Knight, C. G. & Green, N. M.) 225–236
1,3,2-Dioxaphosphorinane 2-oxides, formation of an unstable covalent intermediate during the inhibition by, of the activity of electric-eel acetylcholinesterase (Ashani, Y. & Leader, H.) 781–790
Dissociation constants, molecular, derivation of values for the negative logarithm (p*K*) of, of dibasic acids from dependences on pH (Dixon, H. B. F.) 249–250
Dodecyl sulphate, sodium, effects of, on the adsorption of proteins on *N*-(3-carboxypropionyl)aminodecyl-Sepharose (Yon, R. J. & Simmonds, R. J.) 417–424
Dogfish (*Squalus acanthias*), kinetic studies on liver glutamate dehydrogenase from (Electricwala, A. H. & Dickinson, F. M.) 449–459
Dolphin fish (*Coryphaena hippurus*), mechanism of the reduction by ascorbate of cytochrome *c* from, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648

Egg-white, chicken, isolation and characterization of thiamin-binding protein from, and its interaction with riboflavin-binding protein (Muniyappa, K. & Adiga, P. R.) 887–894
Egg-white, turtle-dove, purification of ovomucoid from, and its activity as a proteinase inhibitor and as a human P_1-blood-group antigen (François-Gerard, C., Gerday, C. & Beeley, J. G.) 679–685
Elastin, additional evidence for the existence of a precursor to tropoelastin involved in the biosynthesis of, in chick aorta (Heng-Khoo, C. S., Rucker, R. B. & Buckingham, K. W.) 559–567
Electric eel (*Electrophorus electricus*), formation of an unstable covalent intermediate during the inhibition by 1,3,2-dioxaphosphorinane 2-oxides of the activity of acetylcholinesterase from (Ashani, Y. & Leader, H.) 781–790
Electrophorus electricus, *see* Electric eel
Enzymes, patterns of apparent co-operativity in a simple random non-equilibrium mechanism for the interaction of substrates, modifiers and, and comparison with equilibrium allosteric models (Whitehead, E. P. & Egmond, M. R.) 631–639
Epidermis, hoof and snout, ox, intraspecies heterogeneity of keratins of (Lee, L. D., Kubilus, J. & Baden, H. P.) 187–196
Erythrocyte 'ghosts', human, effect of the ratio of cholesterol to phospholipid in the membrane of, on cholesterol exchange (Poznansky, M. J. & Czekanski, S.) 989–991
Erythrocytes, ox, binding of zinc ions to the apoenzyme of superoxide dismutase from (Cass, A. E. G., Hill, H. A. O., Bannister, J. V. & Bannister, W. H.) 477–486
Erythrocytes, ox, study by nuclear-magnetic-relaxation measurements of the properties of the cupric ion sites of superoxide dismutase from (Boden, N., Holmes, M. C. & Knowles, P. F.) 303–309
Escherichia coli, determination of the redox potentials of molybdenum and iron–sulphur centres in nitrate reductase from (Vincent, S. P.) 757–759
Escherichia coli, interaction of the *lacZ* β-galactosidase of, with competitive inhibitors and the role of magnesium ions and ionizations in the activity of the enzyme (Loeffler, R. S. T., Sinnott, M. L., Sykes, B. D. & Withers, S. G.) 145–152
Escherichia coli K12, polypeptide-chain stoicheiometry and lipoic acid content of the pyruvate dehydrogenase complex of (Hale, G. & Perham, R. N.) 129–137
Esterase, neurotoxic, brain, rapid aging of, after inhibition by di-isopropyl phosphorofluoridate (Clothier, B. & Johnson, M. K.) 549–558
Estradiol, *see* Oestradiol
Estrogens, *see* Oestrogens
Ethidium bromide, use of, in a rapid and sensitive method for determination of ribonucleic acid in ribosomal preparations (Mackie, C., Bryant, T. H. E. & Mowbray, J.) 977–979
Eye, sheep, use of cross-linking reagents in an investigation of the organization of rhodopsin in the photoreceptor membrane of rod outer segments of retina of (Brett, M. & Findlay, J. B. C.) 215–223

Fatty acids, compositions of glycolipids and, of two lines of cultured dog kidney cells (Hougland, A. E., Gaush, C. R. & Mayberry, W. R.) 311–317
Ferredoxin component, purification and characterization of, of chick kidney 25-hydroxycholecalciferol 1α-hydroxylase (Kulkoski, J. A. & Ghazarian, J. G.) 673–678
Ferrihaemoglobin, human, breakdown of 2,3-bis-(phospho)-D-glycerate catalysed by (El-Yassin, D. I., Fell, D. A., Lloyd, B. B. & Fisher, R. B.) 373–375
Ferritin, spleen, horse, crystallized, removal of cadmium (II) from (Hegenauer, J., Saltman, P. & Hatlen, L.) 693–695
Formaldehyde dehydrogenase, liver, human, steady-state kinetic model for the mechanism of the reaction catalysed by (Uotila, L. & Mannervik, B.) 869–878
N^{10}-Formyltetrahydrofolate dehydrogenase, roles of pyruvate carboxylase and, in rat liver in the toxic effects of excess of dietary histidine (Scrutton, M. C. & Beis, I.) 833–846
Fowl-pest virus, *see* Newcastle-disease virus
α-L-Fucosidase, liver, human, subunit composition of (Alhadeff, J. A. & Andrews-Smith, G. L.) 753–756
Fumarase, heart, pig, use of affinity chromatography for the purification of (Chaudhuri, S. & Thomas, E. W.) 115–119

β-Galactosidase, *lacZ*, *Escherichia coli*, interaction of, with competitive inhibitors and the role of magnesium ions and ionizations in the activity of the enzyme (Loeffler, R. S. T., Sinnott, M. L., Sykes, B. D. & Withers, S. G.) 145–152

1,3-β-Glucanase, endo-, purification and characterization of, secreted into the culture medium by *Candida utilis* C.E.C.T. 1061 (Villa, T. G., Notario, V. & Villanueva, J. R.) 107–114
Glucoamylase/maltase, small-intestine, rat, enzyme activity in partly dissociated fragments of (Flanagan, P. R. & Forstner, G. G.) 487–492
β-Glucosidases, *Botryodiplodia theobromae*, separation and characterization of, and cellulases (Umezurike, G. M.) 9–19
Glucuronides, oestrogen, direct immunoassay of, in female human urine (Baker, T. S., Jennison, K. M. & Kellie, A. E.) 729–738
Glutamate dehydrogenase, liver, dogfish, kinetic studies on (Electricwala, A. H. & Dickinson, F. M.) 449–459
Glutathione, kinetic analysis supporting a steady-state random sequential mechanism for the multiple inhibition by derivatives of, of rat liver glutathione *S*-transferase A (Jakobson, I., Warholm, M. & Mannervik, B.) 861–868
Glutathione peroxidase, liver, guinea-pig, non-selenium-dependent, reduction of lipid hydroperoxides and steroid hydroperoxides catalysed by (Shreve, M. R., Morrissey, P. G. & O'Brien, P. J.) 761–763
Glutathione peroxidase, placental, human, purification and properties of (Awasthi, Y. C., Dao, D. D., Lal, A. K. & Srivastava, S. K.) 471–476
Glutathione, reduced, role of, as an allosteric activator in a steady-state kinetic model for the mechanism of the reaction catalysed by human liver formaldehyde dehydrogenase (Uotila, L. & Mannervik, B.) 869–878
Glutathione *S*-transferase A, liver, rat, kinetic analysis supporting a steady-state random sequential mechanism for the multiple inhibition by glutathione derivatives of (Jakobson, I., Warholm, M. & Mannervik, B.) 861–868
Glutathione *S*-transferase B (ligandin), liver, rat, effects of modification of the thiol groups on the binding and catalytic activities of (Carne, T., Tipping, E. & Ketterer, B.) 433–439
Glycerol phosphate acyltransferase, liver, rat, evidence for the existence of mitochondrial and microsomal isoenzymes of (Nimmo, H. G.) 283–288
Glycine ethyl ester, activation by, of ox spleen cathepsin D (Dionyssiou-Asteriou, A. & Rakitzis, E. T.) 355–356
Glycolipids, compositions of fatty acids and, of two lines of cultured dog kidney cells (Hougland, A. E., Gaush, C. R. & Mayberry, W. R.) 311–317
Glycoprotein, high-molecular-weight, characterization of, isolated from the pulmonary secretions of human patients with alveolar proteinosis (Sahu, S. & Lynn, W. S.) 153–158
Glycosides, hydroxylysine-linked, characterization of, of human complement subcomponent C1q and of various collagens (Shinkai, H. & Yonemasu, K.) 847–852

Haemoglobin, cat, evidence for two chemically distinct sites and a role for histidine and cysteine residues in the binding of triethyltin to (Elliott, B. M., Aldridge, W. N. & Bridges, J. W.) 461–470
Haemoglobin, ferri-, human, breakdown of 2,3-bis-(phospho)-D-glycerate catalysed by (El-Yassin, D. I., Fell, D. A., Lloyd, B. B. & Fisher, R. B.) 373–375
Haemoglobin, human, effects of sodium dodecyl sulphate on the adsorption of, and other proteins on *N*-(3-carboxypropionyl)aminodecyl-Sepharose (Yon, R. J. & Simmonds, R. J.) 417–424
Heart, horse, mechanism of the reduction by ascorbate of cytochrome *c* from, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Heart muscle, rat, selective inactivation by sodium deoxycholate of lactate dehydrogenases from, and other tissues (Lehnert, T. & Berlet, H. H.) 813–818
Heart, ox, evidence for the presence of type-I copper in a product (Compound C_2) of the reaction of oxygen with the mixed-valence state of mitochondrial cytochrome *c* oxidase from (Chance, B., Saronio, C. & Leigh, J. S., Jr.) 931–941
Heart, ox, low-temperature kinetics of the reaction of fully reduced mitochondrial membrane-bound cytochrome *c* oxidase from, with oxygen studied by spectroscopy in the Soret, α-band and near-infrared regions (Clore, G. M. & Chance, E. M.) 613–621
Heart, pig, enzyme-induced inactivation by acetylenic and vinyl analogues of 4-aminobutyrate of aspartate aminotransferase from, and other aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728
Heart, pig, formation of diastereoisomers of methylcitrate catalysed by citrate synthase from, and its relevance to the appearance of methylcitrate in the urine of human patients with propionic acidaemia (Weidman, S. W. & Drysdale, G. R.) 169–174
Heart, pig, generation by deamidation of multiple forms of aspartate aminotransferase from (Williams, J. A. & John, R. A.) 121–127
Heart, pig, mechanism of the formation of the adduct between oxidized nicotinamide–adenine dinucleotide and pyruvate that is bound to lactate dehydrogenase from (Wilton, D. C.) 951–957
Heart, pig, preparation of stable complexes of oxidized nicotinamide–adenine dinucleotide and derivatives with lactate dehydrogenase from, with endogenous catalytic activity (Gacesa, P. & Venn, R. F.) 369–372
Heart, pig, properties of lactate dehydrogenase from, in which the 'essential' cysteine residue 165 has been modified by thioalkylation (Bloxham, D. P., Sharma, R. P. & Wilton, D. C.) 769–780
Heart, pig, use of affinity chromatography for the purification of fumarase from (Chaudhuri, S. & Thomas, E. W.) 115–119
Hen, *see* Chicken
Hepatocytes, rat, isolated, utilization of endogenous and exogenous sources of substrate for the biosynthesis of cholesterol by (Gibbons, G. F. & Pullinger, C. R.) 255–263
Hexosaminidase, *see N*-Acetylhexosaminidase
Histidine, dietary, roles of pyruvate carboxylase and N^{10}-formyltetrahydrofolate dehydrogenase in rat liver in the toxic effects of excess of (Scrutton, M. C. & Beis, I.) 833–846
Histidine residues, evidence for two chemically distinct sites and a role for cysteine residues and, in the binding of triethyltin to cat haemoglobin (Elliott, B. M., Aldridge, W. N. & Bridges, J. W.) 461–470

Histones, chromatin, *Saccharomyces carlsbergensis*, isolation and characterization of (Pastink, A., Berkhout, T. A., Mager, W. H. & Planta, R. J.) 917–923

Homoserine residue, formation of, during the partial non-cleavage of a methionine–cystine bond in human serum albumin and in ox α-lactalbumin (Doyen, N. & Lapresle, C.) 251–254

Housefly (*Musca domestica*), activity of uridine diphosphate glucosyltransferase in microsomal fraction from (Morello, A. & Repetto, Y.) 809–812

Hyaluronate, role of link-protein in the structure of aggregates of, and pig laryngeal cartilage proteoglycans (Hardingham, T. E.) 237–247

Hydrogen, measurement *in vivo* of the evolution of, by *Anabaena cylindrica* that is catalysed by hydrogenase independently of nitrogenase (Daday, A., Lambert, G. R. & Smith, G. D.) 139–144

Hydrogenase, measurement *in vivo* of the evolution of hydrogen by *Anabaena cylindrica* that is catalysed by, independently of nitrogenase (Daday, A., Lambert, G. R. & Smith, G. D.) 139–144

Hydrolases, lysosomal, characterization of, in urine of human patients with mucolipidosis II (I-cell disease) and mucolipidosis III (Kress, C. B. & Miller, A. L.) 409–415

Hydrolases, lysosomal, liver, human, binding by immobilized lectins of (Fiddler, M. B., Ben-Yoseph, Y. & Nadler, H. L.) 175–180

3(17)β-Hydroxy steroid dehydrogenase, *Pseudomonas testosteroni*, 'affinity' chromatography of, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408

3-Hydroxyanhydroretinol, identification of the naturally occurring anhydrovitamin A_2 of *Saccobranchus fossilis* as (Barua, A. B., Verma, K. & Das, S. R.) 791–799

25-Hydroxycholecalciferol 1α-hydroxylase, kidney, chick, purification and characterization of the ferredoxin component of (Kulkoski, J. A. & Ghazarian, J. G.) 673–678

Hydroxylysine residues, characterization of the glycosides linked to, of human complement subcomponent C1q and of various collagens (Shinkai, H. & Yonemasu, K.) 847–852

I-cell disease (mucolipidosis II), characterization of lysosomal hydrolases in urine of human patients with, and mucolipidosis III (Kress, B. C. & Miller, A. L.) 409–415

Intestine, large, rat, multiple forms and kinetic properties of β-*N*-acetylhexosaminidases from mucosa of, and from colonic tumours from animals treated with 1,2-dimethylhydrazine (Mian, N., Herries, D. G., Cowen, D. M. & Batte, E. A.) 319–330

Intestine, small, rabbit, purification and properties of cytosolic aminotripeptidase from the mucosa of (Doumeng, C. & Maroux, S.) 801–808

Intestine, small, rat, enzyme activity in partly dissociated fragments of maltase/glucoamylase from (Flanagan, P. R. & Forstner, G. G.) 487–492

Intestine, small, rat, inhibition by β-2-thienyl-DL-alanine of the activity of phenylalanine hydroxylase in rat liver and kidney and of the transport of phenylalanine by (Wapnir, R. A. & Moak, G. S.) 347–352

Iodoacetamide, reaction of some derivatives of, with rabbit skeletal-muscle creatine kinase (Price, N. C.) 603–612

Iron–sulphur centre, characterization of, of reduced nicotinamide–adenine dinucleotide–acceptor oxidoreductase (component C) of *Methylococcus capsulatus* methane mono-oxygenase complex (Colby, J. & Dalton, H.) 903–908

Iron–sulphur centres, determination of the redox potentials of molybdenum and, in *Escherichia coli* nitrate reductase (Vincent, S. P.) 757–759

Iron–sulphur proteins, properties of, of the *Pseudomonas putida* benzene dioxygenase system (Crutcher, S. E. & Geary, P. J.) 393–400

Kallikrein, submandibular-gland, pig, isolation and properties of (Lemon, M., Fiedler, F., Förg-Brey, B., Hirschauer, C., Leysath, G. & Fritz, H.) 159–168

Keratins, epidermal, hoof and snout, ox, intraspecies heterogeneity of (Lee, L. D., Kubilus, J. & Baden, H. P.) 187–196

Kidney cells, dog, cultured, compositions of glycolipids and fatty acids of two lines of (Hougland, A. E., Gaush, C. R. & Mayberry, W. R.) 311–317

Kidney cells, rat, normal and transformed, activities of diamine oxidase and polyamine oxidase in (Quash, G., Keolouangkhot, T., Gazzolo, L., Ripoll, H. & Saez, S.) 275–282

Kidney, chick, purification and characterization of the ferredoxin component of 25-hydroxycholecalciferol 1α-hydroxylase from (Kulkoski, J. A. & Ghazarian, J. G.) 673–678

Kidney, pig, kinetics and mechanism of action of aldehyde reductase from (Davidson, W. S. & Flynn, T. G.) 595–601

Kidney, rabbit, requirement for negatively charged dispersions of phospholipids for interaction with lipid-depleted sodium-plus-potassium ion-dependent adenosine triphosphatase from (Isern de Caldentey, M. & Wheeler, K. P.) 265–273

Kidney, rat, inhibition by β-2-thienyl-DL-alanine of the activity of phenylalanine hydroxylase in, and liver and of the transport of phenylalanine by the small intestine (Wapnir, R. A. & Moak, G. S.) 347–352

α-Lactalbumin, ox, partial non-cleavage by cyanogen bromide of a methionine–cystine bond in (Doyen, N. & Lapresle, C.) 251–254

β-Lactamase I, *Bacillus cereus*, inactivation by 6β-bromopenicillanic acid of (Knott-Hunziker, V., Orlek, B. S., Sammes, P. G. & Waley, S. G.) 365–367

Lactate dehydrogenase, heart, pig, mechanism of the formation of the adduct between oxidized nicotinamide-adenine dinucleotide and pyruvate that is bound to (Wilton, D. C.) 951–957

Lactate dehydrogenase, heart, pig, preparation of stable complexes of oxidized nicotinamide–adenine dinucleotide and derivatives with, with endogenous catalytic activity (Gacesa, P. & Venn, R. F.) 369–372

Lactate dehydrogenase, heart, pig, properties of, in which the 'essential' cysteine residue 165 has been modified by thioalkylation (Bloxham, D. P., Sharma, R. P. & Wilton, D. C.) 769–780

Lactate dehydrogenases, selective inactivation by sodium deoxycholate of, from various rat tissues (Lehnert, T. & Berlet, H. H.) 813–818

Lactobacillus casei MTX/R, circular-dichroism studies of the binding of ligands to tetrahydrofolate dehydrogenase from (Hood, K., Bayley, P. M. & Roberts, G. C. K.) 425–432

Lactuca scarida, *see* Lettuce

Large intestine, *see* Intestine, large

Larynx, pig, role of link-protein in the structure of aggregates of hyaluronate and proteoglycans from cartilage of (Hardingham, T. E.) 237–247

Leaves, lettuce, purification by ion-exchange chromatography of the chloroplast membrane dicyclohexylcarbodi-imide-binding proteolipid from (Sigrist-Nelson, K. & Azzi, A.) 687–692

Lectins, immobilized, binding by, of human liver lysosomal hydrolases (Fiddler, M. B., Ben-Yoseph, Y. & Nadler, H. L.) 175–180

Legumin, pea-seed, use of immunoaffinity chromatography for the purification of (Casey, R.) 509–520

Lettuce (*Lactuca scarida*) leaves, purification by ion-exchange chromatography of the chloroplast membrane dicyclohexylcarbodi-imide-binding proteolipid from (Sigrist-Nelson, K. & Azzi, A.) 687–692

Ligandin (glutathione *S*-transferase B), liver, rat, effects of modification of the thiol groups on the binding and catalytic activities of (Carne, T., Tipping, E. & Ketterer, B.) 433–439

Light, activation by α-chymotrypsin of silkworm tyrosinase precursor and its relevance to the induction by, of pigmentation (Kuan, K. N., Lee, Y. Y. & Melius, P.) 981–983

Lipid hydroperoxides, reduction of steroid hydroperoxides and, catalysed by a non-selenium-dependent guinea-pig liver glutathione peroxidase (Shreve, M. R., Morrissey, P. G. & O'Brien, P. J.) 761–763

Lipoic acid, polypeptide-chain stoicheiometry and content of, of *Escherichia coli* K12 pyruvate dehydrogenase complex (Hale, G. & Perham, R. N.) 129–137

Liver cells, rat, isolated, utilization of endogenous and exogenous sources of substrate for the biosynthesis of cholesterol by (Gibbons, G. F. & Pullinger, C. R.) 255–263

Liver, chicken, 'affinity' chromatography of oestradiol-17β dehydrogenase from, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)-aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408

Liver, chicken, ligand-induced conformational transitions and secondary structure composition of pyruvate carboxylase from (McGurk, K. S. & Spivey, H. O.) 697–705

Liver, dogfish, kinetic studies on glutamate dehydrogenase from (Electricwala, A. H. & Dickinson, F. M.) 449–459

Liver, frog, synthesis of cytidine monophosphate sialic acids by using cytidine monophosphate acylneuraminate synthase from, immobilized on Sepharose 4B (Corfield, A. P., Schauer, R. & Wember, M.) 1–7

Liver, guinea-pig, reduction of lipid hydroperoxides and steroid hydroperoxides catalysed by a non-selenium-dependent glutathione peroxidase from (Shreve, M. R., Morrissey, P. G. & O'Brien, P. J.) 761–763

Liver, hamster, persistence of methylated base residues in ribonucleic acids of, after administration of dimethylnitrosamine to the animals (Margison, G. P., Margison, J. M. & Montesano, R.) 967–973

Liver, human, binding by immobilized lectins of lysosomal hydrolases from (Fiddler, M. B., Ben-Yoseph, Y. & Nadler, H. L.) 175–180

Liver, human, partial purification of human milk alkaline phosphatase and comparison of its properties with those of the isoenzymes from, and placenta (Hamilton, T. A., Górnicki, S. Z. & Sussman, H. H.) 197–201

Liver, human, steady-state kinetic model for the mechanism of the reaction catalysed by formaldehyde dehydrogenase from (Uotila, L. & Mannervik, B.) 869–878

Liver, human, subunit composition of α-L-fucosidase from (Alhadeff, J. A. & Andrews-Smith, G. L.) 753–756

Liver, mouse, properties of mitochondrial monoamine oxidase from, of obese–hyperglycaemic (*ob*/*ob*) animals (Tong, J. H., Limson-Zamora, M., D'Iorio, A. & Bégin-Heick, N.) 943–949

Liver, rabbit, use of ^{31}P nuclear-magnetic-resonance spectroscopy to study the metabolism of nucleotides by microsomal uridine diphosphate glucuronyltransferase and nucleoside diphosphatase from (Finch, S. A. E., Slater, T. F. & Stier, A.) 925–930

Liver, rat, comparison of endogenous and exogenous ribonucleic acids as primers for the activity of polyuridylate polymerase in ribosomes from (Hayashi, T. T. & MacFarlane, K.) 895–902

Liver, rat, disaggregation of plasma-membrane adenylate cyclase from, during polyacrylamide-gel electrophoresis in mixtures of ionic and non-ionic detergents (Newby, A. C. & Chrambach, A.) 623–630

Liver, rat, effects of modification of the thiol groups on the binding and catalytic activities of ligandin from (Carne, T., Tipping, E. & Ketterer, B.) 433–439

Liver, rat, enzyme-induced inactivation by acetylenic and vinyl analogues of 4-aminobutyrate of ornithine aminotransferase from, and other aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728

Liver, rat, evidence for the existence of isoenzymes of glycerol phosphate acyltransferase in the mitochondrial and microsomal fractions of (Nimmo, H. G.) 283–288

Liver, rat, Gunn, stimulation *in vitro* by alkyl ketones of the activity of defective uridine diphosphate glucuronyltransferase from (Lalani, E.-N. M. A. & Burchell, B.) 993–995

Liver, rat, induction by diamines of inhibition of the activity of ornithine decarboxylase in (Kallio, A., Löfman, M., Pösö, H. & Jänne, J.) 63–69

Liver, rat, inhibition by β-2-thienyl-DL-alanine of the activity of phenylalanine hydroxylase in, and kidney and of the transport of phenylalanine by the small intestine (Wapnir, R. A. & Moak, G. S.) 347–352

Liver, rat, kinetic analysis supporting a steady-state random sequential mechanism for the multiple inhibition by glutathione derivatives of glutathione *S*-transferase A from (Jakobson, I., Warholm, M. & Mannervik, B.) 861–868
Liver, rat, purification and properties of a cytosol protein factor from, that binds and deacylates initiator transfer ribonucleic acid (Pohlreich, P., Kříž, O., Tuháčková, Z., Dušek, Z. & Hradec, J.) 707–719
Liver, rat, roles of pyruvate carboxylase and N^{10}-formyltetrahydrofolate dehydrogenase in, in the toxic effects of excess of dietary histidine (Scrutton, M. C. & Beis, I.) 833–846
Liver, rat, selective inactivation by sodium deoxycholate of lactate dehydrogenases from, and other tissues (Lehnert, T. & Berlet, H. H.) 813–818
Liver, rat, subcellular distribution and substrate specificity of the activities of acetyl-coenzyme A deacylase in (Grigat, K.-P., Koppe, K., Seufert, C.-D. & Söling, H.-D.) 71–79
Liver, sheep, purification of mitochondrial aspartate aminotransferase from, and comparison of its kinetic and other properties with those of the cytosolic isoenzyme (Orlacchio, A., Campos-Cavieres, M., Pashev, I. & Munn, E. A.) 583–593
Lung, human, characterization of a high-molecular-weight glycoprotein isolated from the pulmonary secretions of, of patients with alveolar proteinosis (Sahu, S. & Lynn, W. S.) 153–158
Lysine residues, interaction of dinitrophenyl groups bound to, of ox serum albumin with univalent fragments of anti-dinitrophenyl antibody (Knight, C. G. & Green, N. M.) 225–236
Lysosomes, characterization of hydrolases derived from, in urine of human patients with mucolipidosis II (I-cell disease) and mucolipidosis III (Kress, B. C. & Miller, A. L.) 409–415
Lysosomes, liver, human, binding by immobilized lectins of hydrolases from (Fiddler, M. B., Ben-Yoseph, Y. & Nadler, H. L.) 175–180
Lysyl oxidase, aorta, ox, purification and properties of four species of (Kagan, H. M., Sullivan, K. A., Olsson, T. A., III & Cronlund, A. L.) 203–214

α_2-Macroglobulin, human, identification of distinctive forms of, by using the numerical relationship between the binding of ox trypsin in the α- and β-modes (Topping, R. M. & Craven, A. H.) 501–508
Magnesium ions, interaction of *Escherichia coli lacZ* β-galactosidase with competitive inhibitors and the role of, and ionizations in the activity of the enzyme (Loeffler, R. S. T., Sinnott, M. L., Sykes, B. D. & Withers, S. G.) 145–152
Malate dehydrogenases, bacterial, simple efficient methods for the purification of (Wright, I. P. & Sundaram, T. K.) 441–448
Maltase/glucoamylase, small-intestine, rat, enzyme activity in partly dissociated fragments of (Flanagan, P. R. & Forstner, G. G.) 487–492
Melanin, activation by α-chymotrypsin of silkworm tyrosinase precursor and its relevance to the light-induced formation of (Kuan, K. N., Lee, Y. Y. & Melius, P.) 981–983
Membrane, chloroplast, lettuce-leaf, purification by ion-exchange chromatography of the dicyclohexylcarbodi-imide-binding proteolipid from (Sigrist-Nelson, K. & Azzi, A.) 687–692
Membrane, erythrocyte-'ghost', human, effect of the ratio of cholesterol to phospholipid in, on cholesterol exchange (Poznansky, M. J. & Czekanski, S.) 989–991
Membrane, mitochondrial, heart, ox, low-temperature kinetics of the reaction of fully reduced cytochrome *c* oxidase bound to, with oxygen studied by spectroscopy in the Soret, α-band and near-infrared regions (Clore, G. M. & Chance, E. M.) 613–621
Membrane, photoreceptor, rod-outer-segment, retina, sheep, use of cross-linking reagents in an investigation of the organization of rhodopsin in (Brett, M. & Findlay, J. B. C.) 215–223
Membrane, plasma, kidney, dog, composition of glycolipids of, from two cultured cell lines (Hougland, A. E., Gaush, C. R. & Mayberry, W. R.) 311–317
Membrane, plasma, liver, rat, disaggregation of adenylate cyclase from, during polyacrylamide-gel electrophoresis in mixtures of ionic and non-ionic detergents (Newby, A. C. & Chrambach, A.) 623–630
Menstrual cycle, direct immunoassay of oestrogen glucuronides in female human urine during (Baker, T. S., Jennison, K. M. & Kellie, A. E.) 729–738
Methane mono-oxygenase complex, *Methylococcus capsulatus*, characterization of the iron–sulphur centre of reduced nicotinamide–adenine dinucleotide–acceptor oxidoreductase (component C) of (Colby, J. & Dalton, H.) 903–908
Methane mono-oxygenases, *Methylococcus capsulatus*, *Methylomonas methanica* and *Methylosinus trichosporium*, comparison of the substrate and electron-donor specificities of (Stirling, D. I., Colby, J. & Dalton, H.) 361–364
Methionine–cystine bond, partial non-cleavage by cyanogen bromide of, in human serum albumin and in ox α-lactalbumin (Doyen, N. & Lapresle, C.) 251–254
Methylation, evidence for, of the precursors of rat brain transfer ribonucleic acids during postnatal development (Elahi, E. & Sellinger, O. Z.) 381–384
Methylcitrate, formation of diastereoisomers of, catalysed by pig heart citrate synthase and its relevance to the appearance of methylcitrate in the urine of human patients with propionic acidaemia (Weidman, S. W. & Drysdale, G. R.) 169–174
Methylococcus capsulatus, characterization of the iron-sulphur centre of reduced nicotinamide–adenine dinucleotide–acceptor oxidoreductase (component C) of methane mono-oxygenase complex from (Colby, J. & Dalton, H.) 903–908
Methylococcus capsulatus, comparison of the substrate and electron-donor specificities of methane mono-oxygenases from *Methylomonas methanica*, *Methylosinus trichosporium* and (Stirling, D. I., Colby, J. & Dalton, H.) 361–364
Methylomonas methanica, comparison of the substrate and electron-donor specificities of methane mono-oxygenases from *Methylococcus capsulatus*, *Methylosinus trichosporium* and (Stirling, D. I., Colby, J. & Dalton, H.) 361–364

Methylosinus trichosporium, comparison of the substrate and electron-donor specificities of methane monooxygenases from *Methylococcus capsulatus*, *Methylomonas methanica* and (Stirling, D. I., Colby, J. & Dalton, H.) 361–364
Microsomal fraction, housefly, activity of uridine diphosphate glucosyltransferase in (Morello, A. & Repetto, Y.) 809–812
Microsomal fraction, liver, rabbit, use of ^{31}P nuclear-magnetic-resonance spectroscopy to study the metabolism of nucleotides by uridine diphosphate glucuronyltransferase and nucleoside diphosphatase from (Finch, S. A. E., Slater, T. F. & Stier, A.) 925–930
Microsomal fraction, liver, rat, evidence for the existence of isoenzymes of glycerol phosphate acyltransferase in, and mitochondria (Nimmo, H. G.) 283–288
Milk, human, partial purification of alkaline phosphatase from, and comparison of its properties with those of the placental and liver isoenzymes (Hamilton, T. A., Górnicki, S. Z. & Sussman, H. H.) 197–201
Milk, relationship between the Rapid and Very Rapid electron-paramagnetic-resonance signals given by molybdenum of xanthine oxidase from (Bray, R. C., Gutteridge, S., Stotter, D. A. & Tanner, S. J.) 357–360
Mitochondria, heart, ox, evidence for the presence of type-I copper in a product (Compound C_2) of the reaction of oxygen with the mixed-valence state of cytochrome *c* oxidase from (Chance, B., Saronio, C. & Leigh, J. S., Jr.) 931–941
Mitochondria, heart, ox, low-temperature kinetics of the reaction of fully reduced membrane-bound cytochrome *c* oxidase from, with oxygen studied by spectroscopy in the Soret, α-band and near-infrared regions (Clore, G. M. & Chance, E. M.) 613–621
Mitochondria, liver, rat, evidence for the existence of isoenzymes of glycerol phosphate acyltransferase in, and microsomal fraction (Nimmo, H. G.) 283–288
Mitochondria, liver, rat, properties of monoamine oxidase from, of obese–hyperglycaemic (*ob*/*ob*) animals (Tong, J. H., Limson-Zamora, M., D'Iorio, A. & Bégin-Heick, N.) 943–949
Mitochondria, liver, sheep, purification of aspartate aminotransferase from, and comparison of its kinetic and other properties with those of the cytosolic isoenzyme (Orlacchio, A., Campos-Cavieres, M., Pashev, I. & Munn, E. A.) 583–593
Molybdenum, relationship between the Rapid and Very Rapid electron-paramagnetic-resonance signals given by, of milk xanthine oxidase (Bray, R. C., Gutteridge, S., Stotter, D. A. & Tanner, S. J.) 357–360
Molybdenum, determination of the redox potentials of iron–sulphur centres and, in *Escherichia coli* nitrate reductase (Vincent, S. P.) 757–759
Monoamine oxidase, mitochondrial, liver, rat, properties of, from obese–hyperglycaemic (*ob*/*ob*) animals (Tong, J. H., Limson-Zamora, M., D'Iorio, A. & Bégin-Heick, N.) 943–949
Mucolipidosis II (I-cell disease), characterization of lysosomal hydrolases in urine of human patients with, and mucolipidosis III (Kress, B. C. & Miller, A. L.) 409–415
Mucolipidosis III, characterization of lysosomal hydrolases in urine of human patients with mucolipidosis II (I-cell diseaase) and (Kress, B. C. & Miller, A. L.) 409–415
Mucosa, colonic, rat, multiple forms and kinetic properties of β-*N*-acetylhexosaminidases from, and from colonic tumours from animals treated with 1,2-dimethylhydrazine (Mian, N., Herries, D. G., Cowen, D. M. & Batte, E. A.) 319–330
Mucosa, small-intestinal, rabbit, purification and properties of cytosolic aminotripeptidase from (Doumeng, C. & Maroux, S.) 801–808
Musca domestica, *see* Housefly
Muscle, skeletal and heart, rat, selective inactivation by sodium deoxycholate of lactate dehydrogenases from, and other tissues (Lehnert, T. & Berlet, H. H.) 813–818
Muscle, skeletal, rabbit, reaction of some derivatives of iodoacetamide with creatine kinase from (Price, N. C.) 603–612
Muscle, smooth, rabbit, purification and characterization of troponin C-like proteins (calmodulins) from, and other tissues (Grand, R. J. A., Perry, S. V. & Weeks, R. A.) 521–529
Myelin sheath, brain, ox, isolated, intrinsic fluorescence of preparations of (Crang, A. J. & Rumsby, M. G.) 739–745
Myometrium, uterus, rat, activities of proteinases and proteinase inhibitor in, during pregnancy and involution (Afting, E.-G., Becker, M. L. & Elce, J. S.) 99–106

Nervous system, central, ox, intrinsic fluorescence of preparations of myelin sheath isolated from (Crang, A. J. & Rumsby, M. G.) 739–745
Neurotoxic esterase, *see* Esterase, neurotoxic
Newcastle-disease virus, use of 12-(4-azido-2-nitrophenoxy)stearoylglucosamine as a photoreactive probe for the mapping of the envelope proteins of (Bramhall, J. S., Shiflett, M. A. & Wisnieski, B. J.) 765–768
Nicotinamide–adenine dinucleotide, oxidized, mechanism of the formation of the adduct between pyruvate and, that is bound to pig heart lactate dehydrogenase (Wilton, D. C.) 951–957
Nicotinamide–adenine dinucleotide, oxidized, preparation of stable complexes of, and derivatives with pig heart lactate dehydrogenase with endogenous catalytic activity (Gacesa, P. & Venn, R. F.) 369–372
Nicotinamide–adenine dinucleotide phosphate, reduced, circular-dichroism studies of the binding of, and other ligands to *Lactobacillus casei* MTX/R tetrahydrofolate dehydrogenase (Hood, K., Bayley, P. M. & Roberts, G. C. K.) 425–432
Nicotinamide–adenine dinucleotide (reduced)–acceptor oxidoreductase (component C), characterization of the iron–sulphur centre of, of *Methylococcus capsulatus* methane mono-oxygenase complex (Colby, J. & Dalton, H.) 903–908
Nitrate reductase, *Escherichia coli*, determination of the redox potentials of molybdenum and iron–sulphur centres in (Vincent, S. P.) 757–759
Nitrogenase, measurement *in vivo* of the evolution of hydrogen by *Anabaena cylindrica* that is catalysed by hydrogenase independently of (Daday, A., Lambert, G. R. & Smith, G. D.) 139–144
Non-histone proteins, *see* Proteins, non-histone
Nuclei, brain, rat, characterization of the chromosomal non-histone proteins of the main classes of, fractionated by zonal centrifugation (Tsitilou, S. G., Cox, D., Mathias, A. P. & Ridge, D.) 331–346

Nucleoside diphosphatase, microsomal, liver, rabbit, use of ^{31}P nuclear-magnetic-resonance spectroscopy to study the metabolism of nucleotides by, and uridine diphosphate glucuronyltransferase (Finch, S. A. E., Slater, T. F. & Stier, A.) 925–930
Nucleotide residues, methylated, persistence of, in hamster liver ribonucleic acids after administration of dimethylnitrosamine to the animals (Margison, G. P., Margison, J. M. & Montesano, R.) 967–973
Nucleotides, use of ^{31}P nuclear-magnetic-resonance spectroscopy to study the metabolism of, by rabbit liver microsomal uridine diphosphate glucuronyltransferase and nucleoside diphosphatase (Finch, S. A. E., Slater, T. F. & Stier, A.) 925–930

Oestrogens, direct immunoassay of glucuronides of, in female human urine (Baker, T. S., Jennison, K. M. & Kellie, A. E.) 729–738
Oestradiol-17β dehydrogenase, liver, chicken, 'affinity' chromatography of, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408
Organophosphorus compounds, formation of an unstable covalent intermediate during the inhibition by, of the activity of electric-eel acetylcholinesterase (Ashani, Y. & Leader, H.) 781–790
Organophosphorus compounds, rapid aging of chicken brain neurotoxic esterase after inhibition by (Clothier, B. & Johnson, M. K.) 549–558
Ornithine aminotransferase, liver, rat, enzyme-induced inactivation by acetylenic and vinyl analogues of 4-aminobutyrate of, and other aminotransferases (John, R. A., Jones, E. D. & Fowler, L. J.) 721–728
Ornithine decarboxylase, induction by diamines of the activity of, in rat liver (Kallio, A., Löfman, M., Pösö, H. & Jänne, J.) 63–69
Ovomucoid, egg-white, turtle-dove, purification of, and its activity as a proteinase inhibitor and as a human P_1-blood-group antigen (François-Gerard, C., Gerday, C. & Beeley, J. G.) 679–685
Oxygen, evidence for the presence of type-I copper in a product (Compound C_2) of the reaction of, with the mixed-valence state of ox heart mitochondrial cytochrome *c* oxidase (Chance, B., Saronio, C. & Leigh, J. S., Jr.) 931–941
Oxygen, low-temperature kinetics of the reaction of fully reduced ox heart mitochondrial membrane-bound cytochrome *c* oxidase with, studied by spectroscopy in the Soret, α-band and near-infrared regions (Clore, G. M. & Chance, E. M.) 613–621

Papain, necessary modification to the preparation of, from any high-quality papaya latex and evidence for the structural integrity of the enzyme produced by traditional methods (Baines, B. S. & Brocklehurst, K.) 541–548
Papaya (*Carica papaya*), necessary modification to the preparation of papain from any high-quality latex from, and evidence for the structural integrity of the enzyme produced by traditional methods (Baines, B. S. & Brocklehurst, K.) 541–548
Pea (*Pisum sativum*) seeds, use of immunoaffinity chromatography for the purification of legumin from (Casey, R.) 509–520
Penicillinase (β-lactamase I), *Bacillus cereus*, inactivation by 6β-bromopenicillanic acid of (Knott-Hunziker, V., Orlek, B. S., Sammes, P. G. & Waley, S. G.) 365–367
Peptidyltransferase–DD-carboxypeptidase, exocellular, *Streptomyces* R61, effects of nucleophiles on the breakdown of the benzylpenicilloyl–enzyme complex EI* formed between benzylpenicillin and (Marquet, A., Frère, J.-M., Ghuysen, J.-M. & Loffet, A.) 909–916
pH, derivation of molecular p*K* values of dibasic acids from dependences on (Dixon, H. B. F.) 249–250
pH, effects of calcium ions and, on the intrinsic fluorescence of ox prothrombin fragment 1 (Scott, M. E., Koehler, K. A. & Hiskey, R. G.) 879–886
Phenobarbital, induction by, of the activity of microsomal uridine diphosphate glucosyltransferase in the housefly (Morello, A. & Repetto, Y.) 809–812
Phenol oxidase precursor, silkworm, activation by α-chymotrypsin of, and its relevance to light-induced pigmentation (Kuan, K. N., Lee, Y. Y. & Melius, P.) 981–983
p-(Phenoxypropoxy)aniline, 'affinity' chromatography of steroid-transforming enzymes on columns of, attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408
Phenylalanine hydroxylase, inhibition by β-2-thienyl-DL-alanine of the activity of, in rat liver and kidney and of the transport of phenylalanine by the small intestine (Wapnir, R. A. & Moak, G. S.) 347–352
Phenylalanine, inhibition by β-2-thienyl-DL-alanine of the activity of phenylalanine hydroxylase in rat liver and kidney and of the transport of, by the small intestine (Wapnir, R. A. & Moak, G. S.) 347–352
Phenyl[2-^{3}H]glyoxal, synthesis of, from [2-^{3}H]acetophenone, and its use in the modification of arginine residues in enzymes (Augustus, B. W. & Hutchinson, D. W.) 377–379
Phenylketonuria, limitations of the potential usefulness of β-2-thienyl-DL-alanine as an inducer of, in the rat (Wapnir, R. A. & Moak, G. S.) 347–352
Phosphatase, alkaline, milk, human, partial purification of, and comparison of its properties with those of the placental and liver isoenzymes (Hamilton, T. A., Górnicki, S. Z. & Sussman, H. H.) 197–201
Phospholipid, effect of the ratio of cholesterol to, in human erythrocyte-'ghost' membrane on cholesterol exchange (Poznansky, M. J. & Czekanski, S.) 989–991
Phospholipids, requirement for negatively charged dispersions of, for interaction with lipid-depleted rabbit kidney sodium-plus-potassium ion-dependent adenosine triphosphatase (Isern de Caldentey, M. & Wheeler, K. P.) 265–273
O-Phosphoserine residues, presence of *O*-phosphothreonine and, in non-collagenous proteins and their absence from phosphorylated collagen from chicken bone (Cohen-Solal, L., Lian, J. B., Kossiva, D. & Glimcher, M. J.) 81–98
O-Phosphothreonine residues, presence of *O*-phosphoserine and, in non-collagenous proteins and their absence from phosphorylated collagen from chicken bone (Cohen-Solal, L., Lian, J. B., Kossiva, D. & Glimcher, M. J.) 81–98

Pisum sativum, *see* Pea
Placenta, human, determination of the carbohydrate compositions of the A and B isoenzymes of *β-N*-acetylhexosaminidase from (Freeze, H., Geiger, B. & Miller, A. L.) 749–752
Placenta, human, partial purification of human milk alkaline phosphatase and comparison of its properties with those of the isoenzymes from, and liver (Hamilton, T. A., Górnicki, S. Z. & Sussman, H. H.) 197–201
Placenta, human, purification and properties of glutathione peroxidase from (Awasthi, Y. C., Dao, D. D., Lal, A. K. & Srivastava, S. K.) 471–476
Plasma membrane, *see* Membrane, plasma
Plasma, pig, study by magnetic-resonance and kinetic methods of the properties of cupric ions in benzylamine oxidase from (Barker, R., Boden, N., Cayley, G., Charlton, S. C., Henson, R., Holmes, M. C., Kelly, I. D. & Knowles, P. F.) 289–302
Polyamine oxidase, activities of diamine oxidase and, in normal and transformed rat kidney cells (Quash, G., Keolouangkhot, T., Gazzolo, L., Ripoll, H. & Saez, S.) 275–282
Polyuridylate polymerase, comparison of endogenous and exogenous ribonucleic acids as primers for the activity of, in rat liver ribosomes (Hayashi, T. T. & MacFarlane, K.) 895–902
Pregnancy, activities of proteinases and proteinase inhibitor in rat uterus myometrium during, and involution (Afting, E.-G., Becker, M. L. & Elce, J. S.) 99–106
Pregnancy, direct immunoassay of oestrogen glucuronides in female human urine during (Baker, T. S., Jennison, K. M. & Kellie, A. E.) 729–738
Propionic acidaemia, formation of diastereoisomers of methylcitrate catalysed by pig heart citrate synthase and its relevance to the appearance of methylcitrate in the urine of human patients with (Weidman, S. W. & Drysdale, G. R.) 169–174
Protease, *see* Proteinase
Protein factor, cytosolic, liver, rat, purification and properties of, that binds and deacylates initiator transfer ribonucleic acid (Pohlreich, P., Kříž, O., Tuháčková, Z., Dušek, Z. & Hradec, J.) 707–719
Protein, riboflavin-binding, chicken-egg-white, isolation and characterization of chicken-egg-white thiamin-binding protein and its interaction with (Muniyappa, K. & Adiga, P. R.) 887–894
Protein, thiamin-binding, chicken-egg-white, isolation and characterization of, and its interaction with riboflavin-binding protein (Muniyappa, K. & Adiga, P. R.) 887–894
Protein-polysaccharides, *see* Proteoglycans
Proteinase inhibitor, activities of proteinases and, in rat uterus myometrium during pregnancy and involution (Afting, E.-G., Becker, M. L. & Elce, J. S.) 99–106
Proteinase inhibitor, purification of turtle-dove egg-white ovomucoid and its activity as a human P_1-blood-group antigen and as (François-Gerard, C., Gerday, C. & Beeley, J. G.) 679–685
Proteinase inhibitor, tick, characterization of, with allergenic activity (Willadsen, P. & Riding, G. A.) 41–47
Proteinases, activities of, and proteinase inhibitor in rat uterus myometrium during pregnancy and involution (Afting, E.-G., Becker, M. L. & Elce, J. S.) 99–106
Proteins, anomalous reaction of 4-chloro-7-nitrobenzofurazan with thiol groups of (Nitta, K., Bratcher, S. C. & Kronman, M. J.) 385–392
Proteins, effects of sodium dodecyl sulphate on the adsorption of, on *N*-(3-carboxypropionyl)aminodecyl-Sepharose (Yon, R. J. & Simmonds, R. J.) 417–424
Proteins, envelope, Newcastle-disease-virus, use of 12-(4-azido-2-nitrophenoxy)stearoylglucosamine as a photoreactive probe for the mapping of (Bramhall, J. S., Shiflett, M. A. & Wisnieski, B. J.) 765–768
Proteins, iron–sulphur, properties of, of the *Pseudomonas putida* benzene dioxygenase system (Crutcher, S. E. & Geary, P. J.) 393–400
Proteins, non-collagenous, presence of *O*-phosphoserine and *O*-phosphothreonine residues in, and their absence from phosphorylated collagen from chicken bone (Cohen-Solal, L., Lian, J. B., Kossiva, D. & Glimcher, M. J.) 81–98
Proteins, non-histone, chromosomal, characterization of, of the main classes of rat brain nuclei fractionated by zonal centrifugation (Tsitilou, S. G., Cox, D., Mathias, A. P. & Ridge, D.) 331–346
Proteins, redox, single-site, mechanism of the reduction by ascorbate of (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Proteins, troponin C-like (calmodulins), purification and characterization of, from rabbit smooth muscle and other tissues (Grand, R. J. A., Perry, S. V. & Weeks, R. A.) 521–529
Proteoglycans, cartilage, laryngeal, pig, role of link-protein in the structure of aggregates of hyaluronate and (Hardingham, T. E.) 237–247
Proteoglycans, Smith-degraded, modified procedure for the preparation of, for use as substrates in the assay of the activity of xylosyltransferase in extracts of chick-embryo cartilage (Sandy, J. D.) 569–574
Proteolipid, dicyclohexylcarbodi-imide-binding, membrane, chloroplast, lettuce-leaf, purification by ion-exchange chromatography of (Sigrist-Nelson, K. & Azzi, A.) 687–692
Prothrombin fragment 1, ox, effects of calcium ions and pH on the intrinsic fluorescence of (Scott, M. E., Koehler, K. A. & Hiskey, R. G.) 879–886
Pseudomonas aeruginosa, mechanism of the reduction by ascorbate of cytochrome c_{551} and azurin from, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Pseudomonas aeruginosa N.C.T.C. 6750, some magnetic properties of cytochrome oxidase from (Walsh, T. A., Johnson, M. K., Greenwood, C., Barber, D., Springall, J. P. & Thomson, A. J.) 29–39
Pseudomonas indigofera, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Pseudomonas putida, properties of the iron–sulphur proteins of the benzene dioxygenase system from (Crutcher, S. E. & Geary, P. J.) 393–400
Pseudomonas testosteroni, 'affinity' chromatography of 3(17)*β*-hydroxy steroid dehydrogenase from, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408

Pulmonary secretions, human, characterization of a high-molecular-weight glycoprotein isolated from, of patients with alveolar proteinosis (Sahu, S. & Lynn, W. S.) 153–158
Pyridoxal 5′-phosphate, role of the binding of, in the control of the activity of 5-aminolaevulinate synthetase in *Rhodopseudomonas spheroides* N.C.I.B. 8253 (Davies, R. C. & Neuberger, A.) 661–671
Pyruvate carboxylase, liver, chicken, ligand-induced conformational transitions and secondary structure composition of (McGurk, K. S. & Spivey, H. O.) 697–705
Pyruvate carboxylase, roles of N^{10}-formyltetrahydrofolate dehydrogenase and, in rat liver in the toxic effects of excess of dietary histidine (Scrutton, M. C. & Beis, I.) 833–846
Pyruvate dehydrogenase complex, *Escherichia coli* K12, polypeptide-chain stoicheiometry and lipoic acid content of (Hale, G. & Perham, R. N.) 129–137
Pyruvate, mechanism of the formation of the adduct between oxidized nicotinamide–adenine dinucleotide and, that is bound to pig heart lactate dehydrogenase (Wilton, D. C.) 951–957

Red blood cells, *see* Erythrocytes
Redox potentials, determination of, of molybdenum and iron–sulphur centres in *Escherichia coli* nitrate reductase (Vincent, S. P.) 757–759
Redox proteins, single-site, mechanism of the reduction by ascorabe of (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Restriction endonuclease, *see* Deoxyribonuclease, endo-, restriction
Retina, sheep, use of cross-linking reagents in an investigation of the organization of rhodopsin in the photoreceptor membrane of rod outer segments of (Brett, M. & Findlay, J. B. C.) 215–223
Rhodopseudomonas spheroides N.C.I.B. 8253, purification and properties of the high-activity form of 5-aminolaevulinate synthetase from (Davies, R. C. & Neuberger, A.) 649–659
Rhodopseudomonas spheroides N.C.I.B. 8253, role of the binding of pyridoxal 5′-phosphate in the control of the activity of 5-aminolaevulinate synthetase in (Davies, R. C. & Neuberger, A.) 661–671
Rhodopsin, use of cross-linking reagents in an investigation of the organization of, in the photoreceptor membrane of rod outer segments of sheep retina (Brett, M. & Findlay, J. B. C.) 215–223
Riboflavin-binding protein, chicken-egg-white, isolation and characterization of chicken-egg-white thiamin-binding protein and its interaction with (Muniyappa, K. & Adiga, P. R.) 887–894
Ribonucleic acid polymerase A, deoxyribonucleic acid-dependent, yeast, use of convolution analysis as a mathematical method for the study of transcription by (Cooper, C. S. & Quincey, R. V.) 825–831
Ribonucleic acid polymerase, *Escherichia coli*, anomalous reaction of 4-chloro-7-nitrobenzofuran with thiol groups of, and other proteins (Nitta, K., Bratcher, S. C. & Kronman, M. J.) 385–392
Ribonucleic acid, transfer, initiator, purification and properties of a rat liver cytosolic protein factor that binds and deacylates (Pohlreich, P., Kříž, O., Tuháčková, Z., Dušek, Z. & Hradec, J.) 707–719
Ribonucleic acid, use of ethidium bromide in a rapid and sensitive method for determination of, in ribosomal preparations (Mackie, C., Bryant, T H. E. & Mowbray, J.) 977–979
Ribonucleic acids, endogenous and exogenous, comparison of, as primers for the activity of polyuridylate polymerase in rat liver ribosomes (Hayashi, T. T. & MacFarlane, K.) 895–902
Ribonucleic acids, liver, hamster, persistence of methylated base residues in, after administration of dimethylnitrosamine to the animals (Margison, G. P., Margison, J. M. & Montesano, R.) 967–973
Ribonucleic acids, transfer, brain, rat, evidence for the methylation of precursors of, during postnatal development (Elahi, E. & Sellinger, O. Z.) 381–384
Ribosomes, liver, rat, comparison of endogenous and exogenous ribonucleic acids as primers for the activity of polyuridylate polymerase in (Hayashi, T. T. & MacFarlane, K.) 895–902
Ribosomes, use of ethidium bromide in a rapid and sensitive method for determination of ribonucleic acid in preparations of (Mackie, C., Bryant, T. H. E. & Mowbray, J.) 977–979

Saccharomyces carlsbergensis, isolation and characterization of histones of chromatin from (Pastink, A., Berkhout, T. A., Mager, W. H. & Planta, R. J.) 917–923
Saccharomyces cerevisiae, use of convolution analysis as a mathematical method for the study of transcription by deoxyribonucleic acid-dependent ribonucleic acid polymerase A from (Cooper, C. S. & Quincey, R. V.) 825–831
Saccobranchus fossilis, identification of the naturally occurring anhydrovitamin A_2 of, as 3-hydroxyanhydroretinol (Barua, A. B., Verma, K. & Das, S. R.) 791–799
Salivary gland, submandibular, pig, isolation and properties of kallikrein from (Lemon, M., Fiedler, F., Förg-Brey, B., Hirschauer, C., Leysath, G. & Fritz, H.) 159–168
Seeds, pea, use of immunoaffinity chromatography for the purification of legumin from (Casey, R.) 509–520
Serum albumin, *see* Albumin, serum
Serum, human, four-straight-line model for the characteristics of the binding of ox trypsin by (Topping, R. M. & Seilman, S.) 493–499
Sialic acids, cytidine monophosphate, synthesis of, by using frog liver cytidine monophosphate acylneuraminate synthase immobilized on Sepharose 4B (Corfield, A. P., Schauer, R. & Wember, M.) 1–7
Silkworm (*Bombyx mori*), activation by α-chymotrypsin of tyrosinase precursor from, and its relevance to light-induced pigmentation (Kuan, K. N., Lee, Y. Y. & Melius, P.) 981–983
Skeletal muscle, *see* Muscle, skeletal
Slime mould (*Dictyostelium discoideum*), accumulation of uridine diphosphate glucose pyrophosphorylase by axenically grown amoebae of (Hames, B. D. & Hodson, B. A.) 21–28
Slime mould (*Dictyostelium discoideum*), mechanism of the reduction by ascorbate of cytochrome *c* from, and of other single-site redox proteins (Al-Ayash, A. I. & Wilson, M. T.) 641–648
Small intestine, *see* Intestine, small

Smooth muscle, *see* Muscle, smooth
Sodium deoxycholate, selective inactivation by, of lactate dehydrogenases from various rat tissues (Lehnert, T. & Berlet, H. H.) 813–818
Sodium dodecyl sulphate, effects of, on the adsorption of proteins on *N*-(3-carboxypropionyl)aminodecyl-Sepharose (Yon, R. J. & Simmonds, R. J.) 417–424
Spleen, horse, removal of cadmium(II) from crystallized ferritin from (Hegenauer, J., Saltman, P. & Hatlen, L.) 693–695
Spleen, ox, activation by glycine ethyl ester of cathepsin D from (Dionyssiou-Asteriou, A. & Rakitzis, E. T.) 355–356
Squalus acanthias, *see* Dogfish
Staphylococcus spp. purification, inhibitor studies and *N*-terminal amino acid sequences of variants of chloramphenicol acetyltransferase from (Fitton, J. E. & Shaw, W. V.) 575–582
Steroid hydroperoxides, reduction of lipid hydroperoxides and, catalysed by a non-selenium-dependent guinea-pig liver glutathione reductase (Shreve, M. R., Morrissey, P. G. & O'Brien, P. J.) 761–763
Streptomyces dehydrogenans, 'affinity' chromatography of cortisone reductase from, and other steroid-transforming enzymes on columns of *p*-(phenoxypropoxy)-aniline attached to Sepharose (Renwick, A. G. C., Chambers, S. M. & Willcox, P.) 401–408
Streptomyces R61, effects of nucleophiles on the breakdown of the benzylpenicilloyl–enzyme complex EI* formed between benzylpenicillin and exocellular DD-carboxypeptidase–peptidyltransferase from (Marquet, A., Frère, J.-M., Ghuysen, J.-M. & Loffet, A.) 909–916
Streptopelia risoria, *see* Turtle dove
Subcomponent C1q, complement, human, characterization of the hydroxylysine-linked glycosides of, and of various collagens (Shinkai, H. & Yonemasu, K.) 847–852
Subcomponent C1q, complement, ox, purification and characterization of (Campbell, R. D., Booth, N. A. & Fothergill, J. E.) 531–540
Submandibular gland, pig, isolation and properties of kallikrein from (Lemon, M., Fiedler, F., Förg-Brey, B., Hirschauer, C., Leysath, G. & Fritz, H.) 159–168
Supernatant fraction, *see* Cytosol
Superoxide dismutase, erythrocyte, ox, binding of zinc ions to the apoenzyme of (Cass, A. E. G., Hill, H. A. O., Bannister, J. V. & Bannister, W. H.) 447–486
Superoxide dismutase, erythrocyte, ox, study by nuclear-magnetic-relaxation measurements of the properties of the cupric ion sites of (Boden, N., Holmes, M. C. & Knowles, P. F.) 303–309

Tetrahydrofolate dehydrogenase, *Lactobacillus casei* MTX/R, circular-dichroism studies of the binding of ligands to (Hood, K., Bayley, P. M. & Roberts, G. C. K.) 425–432
Thermoactinomyces sacchari, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Thermomonospora fuscha, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Thermus aquaticus, simple efficient methods for the purification of malate dehydrogenases from, and other bacteria (Wright, I. P. & Sundaram, T. K.) 441–448
Thiamin-binding protein, chicken-egg-white, isolation and characterization of, and its interaction with riboflavin-binding protein (Muniyappa, K. & Adiga, P. R.) 887–894
β-2-Thienyl-DL-alanine, inhibition by, of the activity of rat liver and kidney phenylalanine hydroxylase and of the transport of phenylalanine by the small intestine (Wapnir, R. A. & Moak, G. S.) 347–352
Thioalkylation, properties of pig heart lactate dehydrogenase in which the 'essential' cysteine residue 165 has been modified by (Bloxham, D. P., Sharma, R. P. & Wilton, D. C.) 769–780
Thiol groups, effects of modification of, on the binding and catalytic acitivities of rat liver ligandin (Carne, T., Tipping, E. & Ketterer, B.) 433–439
Thiols, anomalous reaction of 4-chloro-7-nitrobenzofurazan with (Nitta, K., Bratcher, S. C. & Kronman, M. J.) 385–392
Tick (*Boophilus microplus*), characterization of a proteinase inhibitor with allergenic activity from (Willadsen, P. & Riding, G. A.) 41–47
Toxin, extracellular, *Corynebacterium ovis*, purification and partial characterization of (Onon, E. O.) 181–186
Transcription, use of convolution analysis as a mathematical method for the study of, by yeast deoxyribonucleic acid-dependent ribonucleic acid polymerase A (Cooper, C. S. & Quincey, R. V.) 825–831
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Triethyltin, evidence for two chemically distinct sites and a role for histidine and cysteine residues in the binding of, to cat haemoglobin (Elliott, B. M., Aldridge, W. N. & Bridges, J. W.) 461–470
Tropoelastin, aorta, chick, additional evidence for the existence of a precursor to (Heng-Khoo, C. S., Rucker, R. B. & Buckingham, K. W.) 559–567
Troponin C-like proteins (calmodulins), purification and characterization of, from rabbit smooth muscle and other tissues (Grand, R. J. A., Perry, S. V. & Weeks, R. A.) 521–529
Trypsin inhibitor, soya-bean, effects of sodium dodecyl sulphate on the adsorption of, and other proteins on *N*-(3-carboxypropionyl)aminodecyl-Sepharose (Yon, R. J. & Simmonds, R. J.) 417–424
Trypsin inhibitor, tick, characterization of, with allergenic activity (Willadsen, P. & Riding, G. A.) 41–47
Trypsin, ox, four-straight-line model for the characteristics of the binding of, by human blood serum (Topping, R. M. & Seilman, S.) 493–499
Trypsin, ox, identification of distinctive forms of human α_2-macroglobulin by using the numerical relationship between the binding of, in the α- and β-modes (Topping, R. M. & Craven, A. H.) 501–508
Tryptophan residues, intrinsic fluorescence characteristics of tyrosine residues and, of preparations of isolated ox brain myelin sheath (Crang, A. J. & Rumsby, M. G.) 739–745
Tumours, colonic, rat, multiple forms and kinetic properties of *β*-*N*-acetylhexosaminidases from, and from colonic mucosa from animals treated with 1,2-dimethylhydrazine (Mian, N., Herries, D. G., Cowen, D. M. & Batte, E. A.) 319–330

Turtle dove (*Streptopelia risoria*), purification of egg-white ovomucoid from, and its activity as a proteinase inhibitor and as a human P_1-blood-group antigen (François-Gerard, C., Gerday, C. & Beeley, J. G.) 679–685
Tyrosinase precursor, silkworm, activation by α-chymotrypsin of, and its relevance to light-induced pigmentation (Kuan, K. N., Lee, Y. Y. & Melius, P.) 981–983
Tyrosine residues, intrinsic fluorescence characteristics of tryptophan residues and, of preparations of isolated ox brain myelin sheath (Crang, A. J. & Rumsby, M. G.) 739–745

Uridine diphosphate glucose pyrophosphorylase, accumulation of, by axenically grown amoebae of *Dictyostelium discoideum* (Hames, B. D. & Hodson, B. A.) 21–28
Uridine diphosphate glucosyltransferase, activity of, in housefly microsomal fraction (Morello, A. & Repetto, Y.) 809–812
Uridine diphosphate glucuronyltransferase, liver, Gunn-rat, defective, stimulation *in vitro* by alkyl ketones of the activity of (Lalani, E.-N. M. A. & Burchell, B.) 993–995
Uridine diphosphate glucuronyltransferase, microsomal, liver, rabbit, use of ^{31}P nuclear-magnetic-resonance spectroscopy to study the metabolism of nucleotides by, and nucleoside diphosphatase (Finch, S. A. E., Slater, T. F. & Stier, A.) 925–930
Urine, characterization of lysosomal hydrolases in, of human patients with mucolipidosis II (I-cell disease) and mucolipidosis III (Kress, B. C. & Miller, A. L.) 409–415
Urine, human, female, direct radioimmunoassay of oestrogen glucuronides in (Baker, T. S., Jennison, K. M. & Kellie, A. E.) 729–738
Uteroglobulin, uterus, rabbit, determination of the *N*-terminal amino acid sequences of, and its precursor (Atger, M., Mercier, J.-C., Haze, G., Fridlansky, F. & Milgrom, E.) 985–988
Uterus, rabbit, determination of the *N*-terminal amino acid sequences of uteroglobulin and its precursor from (Atger, M., Mercier, J.-C., Haze, G., Fridlansky, F. & Milgrom, E.) 985–988
Uterus, rat, activities of proteinases and proteinase inhibitor in myometrium of, during pregnancy and involution (Afting, E.-G., Becker, M. L. & Elce, J. S.) 99–106

Virus, Newcastle-disease, *see* Newcastle-disease virus
Vitamin A_2, anhydro-, *Saccobranchus fossilis*, naturally occurring, identification of, as 3-hydroxyanhydroretinol (Barua, A. B., Verma, K. & Das, S. R.) 791–799
Vitamin B_1, *see* Thiamin
Vitamin B_2, *see* Riboflavin
Vitamin C, *see* Ascorbate
Vitamin D_3, *see* Cholecalciferol

Xanthine oxidase, milk, relationship between the Rapid and Very Rapid electron-paramagnetic-resonance signals given by molybdenum of (Bray, R. C., Gutteridge, S., Stotter, D. A. & Tanner, S. J.) 357–360
Xylosyltransferase, modified procedure for the preparation of Smith-degraded proteoglycans for use as substrates in the assay of the activity of, in extracts of chick-embryo cartilage (Sandy, J. D.) 569–574

Yeast (*Candida utilis* C.E.C.T. 1061), purification and characterization of endo-1,3-β-glucanase secreted into the culture medium by (Villa, T. G., Notario, V. & Villanueva, J. R.) 107–114
Yeast (*Saccharomyces carlsbergensis*), isolation and characterization of histones of chromatin from (Pastink, A., Berkhout, T. A., Mager, W. H. & Planta, R. J.) 917–923
Yeast (*Saccharomyces cerevisiae*), use of convolution analysis as a mathematical method for the study of transcription by deoxyribonucleic acid-dependent ribonucleic acid polymerase A from (Cooper, C. S. & Quincey, R. V.) 825–831

Zinc ions, binding of, to the apoenzyme of ox erythrocyte superoxide dismutase (Cass, A. E. G., Hill, H. A. O., Bannister, J. V. & Bannister, W. H.) 477–486

JECTS

oteinase
ncy and
ce, J. S.)

irus
naturally
roretinol
799

e Rapid
sonance
R. C.,
) 357-

paration
bstrates
f chick-

ion and
ted into
o, V. &

on and
Pastink,
a, R. J.)

volution
study of
pendent
C. S. &

throcyte
. A. O.,

1979